

Evidence-Based Mental Health



Contents

May 2009 Vol 12 No 2

EBMH Notebook

- 33 Uses and abuses of the Hare Psychopathy Checklist *David Crighton*
- 36 Economic evaluations in evidence based healthcare *Barbara Barrett, Sarah Byford*
- 39 *Et al*


Prognosis


- 41 Review: positive psychological well-being reduces the risk of mortality in both ill and healthy populations
- 42 One-third of the excess death rate in 15–54 year olds in Scotland due to problem drug use
- 43 Early response to treatment predicts remission and recovery at 3 years in people with schizophrenia

Diagnosis


- 44 The 3 item anxiety subscale of the Edinburgh Postpartum Depression Scale may detect postnatal depression as well as the 10 item full scale

Therapeutics

- 45  CBT improves maternal perinatal depression in rural Pakistan
- 46 Brief behavioural intervention for infant sleep problems reduces depression in mothers
- 47 School based intervention improves PTSD symptoms in children affected by political violence
- 48 Family focused therapy shortens recovery time from depression but not mania in adolescents with bipolar disorder
- 49 Review: long term use of antidepressants for bipolar disorder reduces depressive episodes but increases risk of mania
- 50 Switching to aripiprazole from olanzapine leads to weight loss in overweight people with schizophrenia or schizoaffective disorder
- 51 Refractory schizophrenia: adding aripiprazole to clozapine reduces negative but not overall symptoms

- 52 Review: non-pharmacological interventions reduce antipsychotic induced weight gain
- 53 Individual placement and support improves employment of young people with first episode psychosis
- 54 Agomelatine improves symptoms of generalised anxiety disorder
- 55  Adjunctive transdermal oestradiol improves positive symptoms and general psychopathology in women with schizophrenia
- 56 Review: long term psychodynamic psychotherapy improves outcomes in people with complex mental disorders
- 57 Physical activity improves cognitive function in people with memory impairment

Aetiology


- 58 Starting a conventional antipsychotic increases risk of death more than an atypical antipsychotic in elderly people with dementia
- 59  Advanced paternal age increases risk of bipolar disorder in offspring
- 60 Combat exposure increases risk of alcohol misuse in military personnel following deployment
- 61 Small for gestational age boys at increased risk of later psychiatric hospitalisation
- 62 Attention disturbances are associated with low birth weight in an urban community but not a suburban community

Prevalence

- 63 Pervasive behavioural problems are common in children born at less than 26 weeks of gestation
- 64 Low recognition of depression among outpatients by internists in China

Electronic pages

- e1 Purpose and procedure

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org

 equator network

 recycle
When you have finished with this please recycle it