


Evidence-Based Mental Health



Contents

February 2018 Volume 21 Issue 1

Perspective

- 1** Is placebo response in antidepressant trials rising or not? A reanalysis of datasets to conclude this long-lasting controversy
T A Furukawa, A Cipriani, S Leucht, L Z Atkinson, Y Ogawa, N Takeshima, Y Hayasaka, A Chaimani, G Salanti
- 4** The limitations of using randomised controlled trials as a basis for developing treatment guidelines
R Mulder, A B Singh, A Hamilton, P Das, T Outhred, G Morris, D Bassett, B T Baune, M Berk, P Boyce, B Lyndon, G Parker, G S Malhi

Systematic review

- 7** A systematic review of network meta-analyses for pharmacological treatment of common mental disorders
T Williams, D J Stein, J Ipser

Clinical reviews

- 12** Effects of drinking on late-life brain and cognition
A Topiwala, K P Ebmeier
- 16** Engaging families in the management of adolescent self-harm
S Aggarwal, G Patton
- 22** Heart rate variability in bipolar disorder and borderline personality disorder: a clinical review
O Carr, M de Vos, K E A Saunders
- 29** WHO Mental Health Gap Action Programme (mhGAP) Intervention Guide: a systematic review of evidence from low and middle-income countries
R C Keynejad, T Dua, C Barbui, G Thornicroft

Statistics in practice

- 34** Causal inference from experiment and observation
M Zwahlen, G Salanti

Evidence-based case conference

- 39** The role of rTMS for patients with severe PTSD and depression
L D Fryml, G Sahlem, J Fox, E B Short

Further commentaries

Electronic pages

- e1** Multiple indicators of ill health are evident in people with psychosis
D L Foley
- e2** Children with mental health disorders have lower symptoms of depression 3 years after contact with mental health services
P Stallard
- e3** Web-based cognitive behaviour therapy for insomnia shows long-term efficacy in improving chronic insomnia
H Meaklim, D Cunningham
- e4** Multifamily therapy may add to the effectiveness of single-family therapy for adolescents with anorexia nervosa
E K Hughes
- e5** Establishing benefit of emotion-focused, carer-based interventions for adolescents with anorexia nervosa
M L Norris, J Couturier
- e6** L-methylfolate cannot yet be recommended as an add-on treatment in schizophrenia
B Misiak
- e7** NMDA receptor antibodies are found in a small subgroup of patients with first-episode psychosis, but their clinical relevance is unknown
M Schou, S G Saether



This article has been chosen by the Editor to be of special interest or importance and is freely available online.



This journal is a member of and subscribes to the principles of the Committee on Publication Ethics
www.publicationethics.org



When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.





21 (1)

Evid Based Mental Health 2018 21: e3-40

Updated information and services can be found at:
<http://ebmh.bmj.com/content/21/1>

These include:

**Email alerting
service**

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>