



Motivational interviewing is equivalent to more intensive treatment, superior to placebo, and will be tested more widely


Burke BL, Arkowitz H, Menchola M. The efficacy of motivational interviewing: a meta-analysis of controlled clinical trials. *J Consult Clin Psychol* 2003;**71**:843–61.


Q How effective are adaptations of motivational interviewing for changing problem behaviours?

METHODS

 **Design:** Systematic review with meta-analysis.

 **Data sources:** PsycINFO (search date not stated), reference lists of three previous reviews, motivational interviewing website (www.motivationalinterview.org), discussion with motivational interviewing trainers.

 **Study selection and analysis:** Controlled clinical trials were included if the intervention used was based on motivational interviewing principles, was conducted individually and face to face, and used appropriate outcome measures. Effect sizes were calculated using Cohen's d measure.

 **Outcomes:** Effect sizes, combined effect sizes for individual problem areas, longevity of treatment effect, and social impact of treatment.

MAIN RESULTS

Thirty studies met the inclusion criteria (15 targeted alcohol problems, five drug addiction, four diet and exercise habits, two smoking, two HIV risk behaviour, one treatment adherence, and one eating disorders), 11 of which found a statistically significant effect size favouring motivational interviewing. Motivational interviewing was significantly more effective than placebo or no treatment for alcohol or drug abuse, and diet and exercise problems, but not for

smoking or HIV risk behaviour (see table). Motivational interviewing was not significantly better than other active psychotherapeutic treatments for alcohol or drug abuse (see table).

CONCLUSIONS

Motivational interviewing gave moderate effects when compared with placebo or no treatment for alcohol and drug abuse, and problems with diet and exercise. Motivational interviewing was as effective as other psychotherapeutic treatments for alcohol or drug abuse, but had no significant effect on smoking or HIV risk behaviour.

Commentary

Does motivational interviewing present dilemmas for the evidence-based mental health practitioner? Motivational interviewing (MI) is an approach rooted in the Rogerian tradition specifically concerned with the exploration and resolution of ambivalence.¹ It was originally developed as a helpful way to talk to those with drinking problems, and involves elements of directiveness in giving structure and focus to dialogue. It has now been applied to a wide array of health problems, showing considerable promise both as brief treatment and for prevention. The accumulation of evidence for applications has now resulted in this first meta-analysis, which provides a compelling and authoritative evaluation of the existing research literature. This evidence base will expand rapidly with many more trials currently in progress than have been reported to date.

MI is a potentially helpful means of avoiding unproductive confrontations about behaviour and change. Brief applications may be offered as interventions in their own right or by way of preparation for other treatments. Research on dedicated adaptations to the most common mental health problems, however, is currently at a relatively early stage of development compared with applications in the original field of addiction. Careful attention on the part of the evidence-based practitioner to the evolving research literature, including intervention development studies of new adaptations, is thus to be encouraged. This will require attention to evidence beyond that which emerges from trials.

One recent finding is the importance of the quality rather than the quantity of client speech relating to commitment to change,² underlining the specific importance of listening very carefully to clients, and more generally to the construction of an atmosphere of meaningful exchange. Careful eliciting of the values, assumptions, fears, and hopes of the patient is fundamental to both MI and to evidence-based mental health practice, as is skilful interpretation and application of evidence in the interests of the client.³

Jim McCambridge
Wellcome Trust Health Services Research Fellow,
National Addiction Centre, London, UK

- 1 Miller WR, Rollnick S. *Motivational interviewing: preparing people for change*, 2nd ed. New York: Guilford Press, 2002.
- 2 Amrhein PC, Miller WR, Yahne CE, et al. Client commitment language during motivational interviewing predicts drug use outcomes. *J Consult Clin Psychol* 2003;**71**:862–78.
- 3 Szatmari P. The art of evidence-based child psychiatry. *Evid Based Ment Health* 2003;**6**:99–100.

Table Effect sizes for motivational interviewing according to target behaviour (95% confidence intervals)

Target behaviour	Motivational interviewing v placebo or no treatment	Motivational interviewing v active treatments
Alcohol abuse		
Blood alcohol concentration	0.53 (0.20 to 0.86)	–
Standard ethanol content	0.25 (0.13 to 0.37)	0.09 (–0.04 to 0.23)
Drug abuse	0.56 (0.31 to 0.82)	–0.01 (–0.25 to 0.25)
Diet and exercise habits	0.53 (0.32 to 0.62)	–
Smoking	0.11 (–0.05 to 0.27)	–
HIV risk	0.01 (–0.29 to 0.31)	–

For correspondence: B Burke, Department of Psychology, Fort Lewis College, Durango, Colorado, USA; burke-b@fortlewis.edu

Sources of funding: not specified.

EBMH

Motivational interviewing is equivalent to more intensive treatment, superior to placebo, and will be tested more widely

Evid Based Mental Health 2004 7: 52
doi: 10.1136/ebmh.7.2.52

Updated information and services can be found at:
<http://ebmh.bmj.com/content/7/2/52>

References

These include:

This article cites 3 articles, 1 of which you can access for free at:
<http://ebmh.bmj.com/content/7/2/52#ref-list-1>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Substance dependence](#) (407)
[Neurology](#) (1070)
[Clinical trials \(epidemiology\)](#) (992)
[Epidemiology](#) (1570)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>