

Evidence-Based Mental Health

Contents

Purpose and procedure	58	Cognitive behavioural therapy has short term but not long term benefits in people with residual symptoms of depression	75
EBMH Notebook			
Rethinking evidence-based practice for children's mental health	60	Improving quality of primary care reduces depression and improves quality of life in adolescents.....	76
bmjupdates ⁺ , a new FREE service for evidence-based clinical practice.....	62	Review: cognitive behaviour therapy is beneficial in children with anxiety disorders.....	77
Other articles noted	64	Adding mixed amphetamine salts to divalproex sodium improves ADHD symptoms in children with bipolar disorder and comorbid ADHD	78
Prognosis			
Functional impairment recovers after episodes of major depression	65	Valproate reduces alcohol consumption in people with comorbid alcohol dependency and bipolar disorder....	79
Review: depression may increase mortality in coronary heart disease	66	Lithium may reduce gambling severity in pathological gamblers with bipolar disorder	80
Review: after myocardial infarction, depression and poor prognosis are associated.....	67	Adding a nurse-based intervention programme to usual care improves manic symptoms in people with bipolar disorder.....	81
People diagnosed with schizophrenia retain the diagnosis long term	68	Glycine and D-serine improve the negative symptoms of schizophrenia.....	82
Diagnosis			
The Screening Tool for Autism in Two Year Olds can identify children at risk of autism.....	69	Review: pharmacotherapy provides minimal improvements in the neuropsychiatric symptoms of dementia	83
Middelheim Frontality Score may be useful for differentiating between Alzheimer's disease and frontotemporal dementia.....	70	Review: short term psychotherapy is an effective treatment for psychiatric disorders	84
Therapeutics		Aetiology	
Review: hypericum extracts are safer and lead to fewer adverse effects than older standard antidepressants, but have similar tolerability to SSRIs.....	71	Review: migrants are at increased risk of developing schizophrenia.....	85
Metyrapone is an effective adjuvant treatment for major depression	72	Low intelligence increases risk of suicide	86
Bright light treatment an effective addition to antidepressants for non-seasonal depression.....	73	Cannabis use increases the risk of young people developing psychotic symptoms, particularly if already predisposed.....	87
Prophylactic mirtazapine may help to prevent post-stroke depression in people with good cognitive function	74	Prevalence	
		Intoxicant use increased among Norwegian adolescents between 1992 and 2002, and attempted suicides increased among girls.....	88



8 (3)

Evid Based Mental Health 2005 8: 58-88

Updated information and services can be found at:
<http://ebmh.bmj.com/content/8/3.citation>

These include:

**Email alerting
service**

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>