

Evidence-Based Mental Health

Contents

Purpose and procedure	2	Diagnosis		
EBMH notebook	4	Questionnaire for eating disorder diagnoses had good sensitivity in a clinical and non-clinical sample of women	22	
Therapeutics		Prognosis		
Caregiver training delayed admission of patients with dementia to nursing homes	9	Child and maternal psychological status was predictive of psychiatric disorders in youths with IDDM	23	
A “critical time” intervention reduced homelessness in inner city men with psychiatric problems	10	Comorbid OCD, higher score on the MFQ, and older age at presentation predicted persistence of major depression in children and adolescents	24	
Child and parent training sessions led to improved child behaviour in child conduct disorder	11	<i>DSM-III-R</i> and <i>ICD-10</i> diagnoses of schizophrenia had higher predictive validity than that of <i>ICD-9</i> and CATEGO S+ diagnoses	26	
Cognitive behavioural therapy reduced symptoms in patients with obsessive-compulsive disorders with no overt compulsive rituals	12	Review: risk for suicide is increased for most mental disorders where patients require treatment in a hospital setting	27	
Review: exposure with response prevention and serotonergic medication reduce symptoms of obsessive-compulsive disorder	13	Aetiology		
Cognitive behaviour therapy improved functional impairment and fatigue in patients with chronic fatigue syndrome	14	Purchase of a handgun was associated with an increased risk of suicide or death by homicide	28	
Review: risperidone produces more clinical improvement and causes fewer extrapyramidal side effects in schizophrenia	15	Comorbid anxiety and depression in childhood was associated with poorer psychosocial functioning in early adulthood	29	
Sertindole controlled symptoms of schizophrenia without motor side effects	16	Antidepressant exposure before birth was not associated with language, physical, or behavioural development	30	
Review: clozapine reduces relapse and symptoms compared with typical neuroleptic drugs in schizophrenia	17	Psychiatric sickness absence was affected by social support at home and at work	31	
Review: benzodiazepines are more effective than neuroleptics in reducing delirium and seizures in alcohol withdrawal	18	Economics		
Brief advice reduced drinking in non-dependent problem drinkers	19	Review: psychotherapy reduces a variety of costs when used to treat patients with the most severe psychiatric disorders	32	
High dose levomethadyl acetate hydrochloride reduced heroin use in patients dependent on opiates	20	Glossary		Inside back cover
Review: discontinuation rates are the same for serotonin specific reuptake inhibitors and newer tricyclic and heterocyclic antidepressants	21			