

# Evidence-Based Mental Health

## Contents

Purpose and procedure .....	2	Diagnosis		
EBMH notebook .....	4	Questionnaire for eating disorder diagnoses had good sensitivity in a clinical and non-clinical sample of women .....	22	
<b>Therapeutics</b>		<b>Prognosis</b>		
Caregiver training delayed admission of patients with dementia to nursing homes .....	9	Child and maternal psychological status was predictive of psychiatric disorders in youths with IDDM .....	23	
A “critical time” intervention reduced homelessness in inner city men with psychiatric problems .....	10	Comorbid OCD, higher score on the MFQ, and older age at presentation predicted persistence of major depression in children and adolescents .....	24	
Child and parent training sessions led to improved child behaviour in child conduct disorder .....	11	<i>DSM-III-R</i> and <i>ICD-10</i> diagnoses of schizophrenia had higher predictive validity than that of <i>ICD-9</i> and CATEGO S+ diagnoses .....	26	
Cognitive behavioural therapy reduced symptoms in patients with obsessive-compulsive disorders with no overt compulsive rituals .....	12	Review: risk for suicide is increased for most mental disorders where patients require treatment in a hospital setting .....	27	
Review: exposure with response prevention and serotonergic medication reduce symptoms of obsessive-compulsive disorder .....	13	<b>Aetiology</b>		
Cognitive behaviour therapy improved functional impairment and fatigue in patients with chronic fatigue syndrome .....	14	Purchase of a handgun was associated with an increased risk of suicide or death by homicide .....	28	
Review: risperidone produces more clinical improvement and causes fewer extrapyramidal side effects in schizophrenia .....	15	Comorbid anxiety and depression in childhood was associated with poorer psychosocial functioning in early adulthood .....	29	
Sertindole controlled symptoms of schizophrenia without motor side effects .....	16	Antidepressant exposure before birth was not associated with language, physical, or behavioural development .....	30	
Review: clozapine reduces relapse and symptoms compared with typical neuroleptic drugs in schizophrenia .....	17	Psychiatric sickness absence was affected by social support at home and at work .....	31	
Review: benzodiazepines are more effective than neuroleptics in reducing delirium and seizures in alcohol withdrawal .....	18	<b>Economics</b>		
Brief advice reduced drinking in non-dependent problem drinkers .....	19	Review: psychotherapy reduces a variety of costs when used to treat patients with the most severe psychiatric disorders .....	32	
High dose levomethadyl acetate hydrochloride reduced heroin use in patients dependent on opiates .....	20	<b>Glossary</b> .....		Inside back cover
Review: discontinuation rates are the same for serotonin specific reuptake inhibitors and newer tricyclic and heterocyclic antidepressants .....	21			