

Evidence-Based Mental Health

Contents

Purpose and procedure	34	An assertive community treatment programme led to improved housing stability in homeless people with severe mental illness.....	54
Other articles noted.....	36	Olanzapine produced a higher clinical response rate than risperidone in schizophrenia.....	55
EBMH notebook	37	Review: zuclopenthixol is not more effective than other neuroleptic drugs in the acute treatment of schizophrenia	56
Therapeutics			
Fluoxetine reduced depressive symptoms and alcohol consumption in patients with comorbid major depression and alcohol dependence	41	Diagnosis	
Reinforcing behaviours other than abstinence was effective in reducing drug use.....	42	Center for Epidemiologic Studies Depression Scale had high sensitivity and specificity for major depression in older adults.....	57
Cognitive behavioural therapy improved symptoms in children with anxiety disorders	43	Aetiology	
Exposure and response prevention reduced obsessive compulsive disorder symptoms.....	44	Review: childhood sexual abuse is associated with bulimia nervosa especially when there is psychiatric comorbidity	58
Pimozide was superior to haloperidol for children and adolescents with Tourette's disorder and caused fewer side effects	45	Low family cohesion was associated with the incidence of major depressive disorder in adolescents ...	59
Buprenorphine and methadone were equally effective in reducing opiate or cocaine use.....	46	Review: onset of schizophrenia before 22 years of age is associated with abnormal presentation at birth and complicated caesarean birth	60
Alprazolam had short term anxiolytic effectiveness but increased physiological activation in flight phobia	47	Prognosis	
Review: some psychosocial interventions can help children and families cope with chronic health conditions.....	48	Review: few elderly inpatients who are depressed improve	61
Fluoxetine decreased depressive symptoms in children and adolescents with non-psychotic major depressive disorder.....	49	Quality improvement	
Review: selective serotonin reuptake inhibitors differ from tricyclic antidepressants in adverse events.....	50	Review: community mental health team management for adults with severe mental illnesses increases satisfaction with care.....	62
Selective serotonin reuptake inhibitors were less cost effective than initial treatment with tricyclic antidepressants.....	51	Economics	
Home visitation improved some maternal and child outcomes.....	52	A programme of methadone maintenance plus moderate counselling services was cost effective.....	63
Home visits by nurses reduced child abuse and neglect	53	Risperidone and clozapine were more cost effective than haloperidol and chlorpromazine in patients with schizophrenia	64
		Glossary	Inside back cover