

Evidence-Based Mental Health

Contents

Purpose and procedure	98	Diagnosis	
Other articles noted	100	Chinese version of the Edinburgh Postnatal Depression Scale was useful for detecting postnatal depression	120
EBMH notebook	101	<i>DSM-IV</i> criteria were reliable and accurate in differentiating pervasive developmental disorder (PDD) from non-PDD and autism from Asperger's disorder	121
Therapeutics		Prognosis	
Review: cognitive behavioural therapy is effective for anger problems	105	Psychiatric problems at baseline in children with hemiplegia predicted psychiatric problems at 4 years ...	122
Integration of medical and social services for elderly people in the community reduced costs and use of health services	106	Review: individual variables have low power to predict recidivism with sex offenders	123
Treatment of severe and persistent mental illness in a community residential alternative was effective	107	15 years of follow up of non-affective functional psychotic disorders showed a high risk of relapse, suicide, and chronicity	124
Sildenafil improved sexual function in erectile dysfunction	108	Aetiology	
A home based family intervention for children who had deliberately poisoned themselves did not reduce suicidal ideation	109	Earlier onset and increased risk of Alzheimer's disease in Down's syndrome was associated with sex and apolipoprotein E genotype	125
Review: cognitive behaviour therapy facilitates recovery from depressive disorders in children and adolescents ...	110	The apolipoprotein E ϵ 4 allele was associated with Alzheimer's disease in white but not in African-American or Hispanic people	126
Review: antidepressant drugs are effective in dysthymia ..	111	Economics	
Personal therapy reduced adverse outcomes in patients with schizophrenia living with family but increased psychotic relapse rates in those living independently; personal therapy improved social adjustment but increased anxiety	112	Collaborative care improved the cost effectiveness of treatment for major depression in primary care	127
Exposure or cognitive restructuring were more effective than relaxation in treating post-traumatic stress disorder	114	Assertive community treatment for high users of inpatient services had different outcomes at different types of hospitals	128
Review: assertive community treatment is an effective alternative in severe mental disorders	115	Subject index	129
Metrifonate was effective in mild to moderate Alzheimer's disease	116	Author index	134
Review: benzodiazepines and zolpidem are effective for chronic insomnia	117	Contents	
Review: a brief psychological intervention (debriefing) is ineffective in preventing post-traumatic stress disorder	118	Glossary	Inside back cover
Review: oestrogen therapy after menopause may improve cognitive performance and may reduce the risk of developing dementia	119		