

# Evidence-Based Mental Health

## Contents

<b>Purpose and procedure</b> .....	2	Psychiatric telephone contact following emergency department discharge reduces suicide re-attempts in people originally admitted for attempted suicide .....	19
<b>EBMH notebook</b>			
All the evidence that's fit to print .....	4	Review: antidepressant use increases the risk of suicidal behaviour and ideation in children .....	20
Evidence-based psychopharmacology: an agenda for the future .....	4	Interventions targeting joint attention and symbolic play can improve aspects of these skills in young children with autism .....	21
<b><i>Et al.</i></b> .....	7	Bibliotherapy is more effective than waiting list for reducing childhood anxiety disorder, but not as effective as group cognitive behavioural therapy .....	22
<b>Diagnosis</b>		Pregabalin and venlafaxine improve symptoms of generalised anxiety disorder .....	23
Parental report version of the Mood Disorder Questionnaire for adolescents has good sensitivity and specificity for diagnosing bipolar disorder in psychiatric outpatient clinics .....	9	Review: antipsychotic augmentation is effective for people with obsessive-compulsive disorder who are receiving, but not responding to, serotonin reuptake inhibitor monotherapy .....	24
<b>Therapeutics</b>		Group cognitive behavioural therapy increases complete remission of obsessive-compulsive disorder symptoms compared with sertraline, but not response .....	25
Review: antidepressant plus antipsychotic increases clinical response rates in psychotic depression .....	10	Light therapy and fluoxetine similarly effective for improving seasonal affective disorder .....	26
Review: atypical antipsychotics are effective for treating bipolar mania .....	11	Homocysteine-lowering vitamin B supplements do not improve cognitive performance in healthy older adults after two years .....	27
Combined olanzapine plus fluoxetine modestly improves symptoms of acute bipolar I depression compared to lamotrigine .....	12	<b>Prevalence</b>	
Continuing aripiprazole after stabilisation of a manic or mixed episode of bipolar I disorder delays relapse .....	13	About 1% of children in the South Thames region have an autistic spectrum disorder .....	28
Review: aripiprazole for schizophrenia—more high quality research required .....	14	Learning disabilities are the most commonly diagnosed emotional, developmental or behavioural problem of US children .....	29
Risperidone improves symptoms in people who are hospitalised during an acute exacerbation of schizophrenia .....	15	Lifetime prevalence of self-injurious behaviour among US college students is 17% .....	30
Mirtazapine and nortriptyline similarly effective third-line treatments for depression .....	16	Increased prevalence of post-traumatic stress disorder, anxiety and depression in displaced tsunami survivors from southern Thailand .....	31
Higher dose venlafaxine may improve response compared with standard doses in people with major depressive disorder and intolerance or non-responsiveness to SSRIs .....	17	Intermittent explosive disorder is common, has an early age of onset and is associated with the development of other mental disorders in the US population .....	32
Dialectic behaviour therapy reduces suicide attempts compared with non-behavioural psychotherapy in women with borderline personality disorder .....	18		