

Evidence-Based Mental Health

Contents

Purpose and procedure	2	Psychiatric telephone contact following emergency department discharge reduces suicide re-attempts in people originally admitted for attempted suicide	19
EBMH notebook		Review: antidepressant use increases the risk of suicidal behaviour and ideation in children	20
All the evidence that's fit to print	4	Interventions targeting joint attention and symbolic play can improve aspects of these skills in young children with autism	21
Evidence-based psychopharmacology: an agenda for the future	4	Bibliotherapy is more effective than waiting list for reducing childhood anxiety disorder, but not as effective as group cognitive behavioural therapy	22
<i>Et al.</i>	7	Pregabalin and venlafaxine improve symptoms of generalised anxiety disorder	23
Diagnosis		Review: antipsychotic augmentation is effective for people with obsessive-compulsive disorder who are receiving, but not responding to, serotonin reuptake inhibitor monotherapy	24
Parental report version of the Mood Disorder Questionnaire for adolescents has good sensitivity and specificity for diagnosing bipolar disorder in psychiatric outpatient clinics	9	Group cognitive behavioural therapy increases complete remission of obsessive-compulsive disorder symptoms compared with sertraline, but not response	25
Therapeutics		Light therapy and fluoxetine similarly effective for improving seasonal affective disorder	26
Review: antidepressant plus antipsychotic increases clinical response rates in psychotic depression	10	Homocysteine-lowering vitamin B supplements do not improve cognitive performance in healthy older adults after two years	27
Review: atypical antipsychotics are effective for treating bipolar mania	11	Prevalence	
Combined olanzapine plus fluoxetine modestly improves symptoms of acute bipolar I depression compared to lamotrigine	12	About 1% of children in the South Thames region have an autistic spectrum disorder	28
Continuing aripiprazole after stabilisation of a manic or mixed episode of bipolar I disorder delays relapse	13	Learning disabilities are the most commonly diagnosed emotional, developmental or behavioural problem of US children	29
Review: aripiprazole for schizophrenia—more high quality research required	14	Lifetime prevalence of self-injurious behaviour among US college students is 17%	30
Risperidone improves symptoms in people who are hospitalised during an acute exacerbation of schizophrenia	15	Increased prevalence of post-traumatic stress disorder, anxiety and depression in displaced tsunami survivors from southern Thailand	31
Mirtazapine and nortriptyline similarly effective third-line treatments for depression	16	Intermittent explosive disorder is common, has an early age of onset and is associated with the development of other mental disorders in the US population	32
Higher dose venlafaxine may improve response compared with standard doses in people with major depressive disorder and intolerance or non-responsiveness to SSRIs	17		
Dialectic behaviour therapy reduces suicide attempts compared with non-behavioural psychotherapy in women with borderline personality disorder	18		