

Evidence-Based Mental Health

Contents

EBMH notebook

p Value fetishism and use of the Bonferroni adjustment	34	IMPACT collaborative care programme reduces suicide ideation in depressed older adults	51
Experimental interventions with sex offenders: a brief review of their efficacy	35	Review: selective and indicated programmes are more effective than universal programmes for preventing depression in children and adolescents at 6 month follow-up	52
Intervention following deliberate self-harm: enough evidence to act?	37	Review: evidence-based treatments are more effective than usual care for youths with psychological problems and maladaptive behaviour	53
Et al	39	Review: effects of behavioural parent training and cognitive-behavioural therapy on antisocial behaviour in children remain unclear	54
Prognosis		Brief cognitive-behavioural therapy reduces hospital readmission but does not reduce overall symptoms of schizophrenia	55
Remission one year after emergency admission is higher for substance-induced psychosis than primary psychosis	41	Quetiapine is no less effective than risperidone for reducing symptoms of schizophrenia	56
Significant and increasing depressive symptoms are associated with increased risk of new cardiac events in people with a heart attack in the previous year	42	Clozapine improves symptoms but not quality of life compared with other second-generation antipsychotics in people with treatment resistant schizophrenia	57
Phobic anxiety and depression increase the risk of ventricular arrhythmias in people with coronary artery disease	43	Time to discontinuation similar for atypical antipsychotic drugs and placebo in people with Alzheimer's disease	58
Diagnosis		Oxcarbazepine is no more effective than placebo for reducing manic symptoms of bipolar I disorder in children and adolescents	59
The Child Trauma Screening Questionnaire predicts PTSD onset 6 months after traumatic accident better than the Children's Impact of Events Scale	44	Methylphenidate may improve symptoms but does not increase response compared with placebo in preschool children with ADHD	60
Therapeutics		Aetiology	
Escitalopram 10 mg daily is more effective than paroxetine and placebo for generalised anxiety disorder	45	Cannabis use increases risk of developing symptoms of mania	61
Review: little evidence to support dose escalation of selective serotonin reuptake inhibitors in non-responders	46	Adolescent suicidal ideation predicts suicidal behaviour and psychopathological problems at age 30	62
Decision support for primary care clinicians improves process of care but not symptoms in people with depression	47	Prevalence	
Review: enhanced primary care programmes are effective for people with depression but are associated with increased healthcare costs	48	Bipolar I disorder is common in first-degree relatives of children with prepubertal and early adolescent bipolar I disorder	63
Benefits of bright light in non-seasonal major depression not sustained after discontinuation	49	Prevalence of non-fatal suicide attempts in black Americans is about 4%	64
Review: pharmacotherapy and psychotherapy similarly effective for depression in older adults	50		