

Contents

EBMH notebook

Is EBMH too PC?.....	66	Combined fluoxetine plus cognitive behavioural therapy is more effective than monotherapy or placebo for adolescents with depression	84
Appraising the evidence: what is selection bias?.....	67	Cognitive behavioural therapy plus fluoxetine offers some safety advantages over fluoxetine alone in adolescents with depression	85
Arts therapies for people with schizophrenia: an emerging evidence base.....	69	The Webster–Stratton “Incredible Years” parent training programme reduces conduct problems in children.....	86
Et al	71	Cognitive behavioural therapy increases response in people with hypochondriasis.....	87
Prognosis		Dihydrocodeine as effective as methadone for maintenance of treatment for opiate dependence?.....	88
Predictors of symptomatic remission in people with schizophrenia identified.....	73	Aetiology	
Therapeutics		Review: Antidepressants may increase risk of self-harm or suicidal behaviour in children and adolescents	89
Haloperidol, risperidone and olanzapine are similarly effective for first-episode non-affective psychosis but have differing side effects.....	74	Antidepressants increase risk of suicide attempt, but not completed suicide, in people with previous suicide attempts	90
Adherence therapy does not improve quality of life in people with schizophrenia	75	Perinatal factors affect risk of suicide	91
Haloperidol, olanzapine and risperidone reduce treatment failure compared to aripiprazole, quetiapine and ziprasidone in acute schizophrenia	76	Head injury slightly increases risk of non-affective psychosis but not schizophrenia	92
Music therapy improves symptoms in adults hospitalised with schizophrenia	77	Prevalence	
Repetitive transcranial magnetic stimulation is not as effective as electroconvulsive therapy for major depression	78	Obsessive-compulsive disorder is uncommon but associated with high levels of comorbid neuroses, impaired function and increased suicidal acts in people in the UK.....	93
No difference in depression relapse rates between maintenance electroconvulsive therapy and lithium plus nortriptyline	79	Prevalence of clinically diagnosed mental ill-health in adults with intellectual disabilities is around 40%	94
Citalopram, but not interpersonal psychotherapy, improves major depression in people with coronary artery disease.....	80	Ethnic minority groups, particularly African-Caribbean and Black African groups, are at increased risk of psychosis in the UK.....	95
Review: Response to SSRIs in unipolar depression occurs in the first week of treatment.....	81	Lifetime prevalence of psychotic disorders in Finland is 3.1%.....	96
Quetiapine more effective than placebo for depression in bipolar I and II disorder	82	Electronic pages	
Review: Antipsychotic plus mood stabiliser co-therapy is more effective than mood stabiliser mono-therapy at reducing acute bipolar mania.....	83	Purpose and procedure.....	e1