


Evidence-Based Mental Health



Contents

February 2008 Vol 11 No 1

Purpose and procedure

- 1 Purpose and procedure

EBMH Notebook

- 3 A different look
S Reid
- 3 Missing data and the trouble with LOCF
DL Streiner
- 5 Cognitive-behavioural therapies: achievements and challenges
BA Gaudiano
- 7 Et al

Prognosis

- 9 Major psychiatric disorders increase risk of mortality

- 10 About a third of young women are estimated to fully recover from social phobia within 18 months

Diagnosis

- 11 The dementia screening questionnaire for individuals with intellectual disabilities has high sensitivity and specificity in adults with Down's syndrome

Therapeutics

- 12 Cognitive behavioural therapy and interpersonal therapy comparably effective for major depression
- 13 Adding cognitive behavioural therapy to SSRIs does not improve outcomes in adolescents with major depression
- 14 Review: Atypical antipsychotics are effective adjuncts for treatment resistant depression but increase discontinuation due to adverse effects

- 15 Adding aripiprazole improves major depressive disorder following incomplete response to antidepressants alone
- 16 Managed depression care reduces mortality in older adults with major depression
- 17 Group interpersonal therapy reduces depression in adolescent survivors of war


- 18 Cognitive remediation therapy produces moderate improvements in working memory in people with schizophrenia
- 19 Review: Adding Chinese herbal medicine to antipsychotics may improve some outcomes in schizophrenia, but more high quality trials are needed
- 20 Intensive family-based CBT similarly effective to weekly CBT for paediatric obsessive compulsive disorder
- 21 Review: More high quality studies needed to determine the effects on patient outcomes of psychiatric care guideline implementation
- 22 Brief personalised motivational interviewing reduces frequency of marijuana use in regular users ambivalent to change
- 23 The internet-based *My Body, My Life: Body Image Program* for adolescent girls improves body image and disordered eating
- 24 Structured outpatient treatments improve some aspects of borderline personality disorder
- 25 Cognitive therapy improves post-traumatic stress disorder associated with civil conflict in Northern Ireland

Aetiology

- 26 Early childhood factors increase risk of post-traumatic stress disorder
- 27 Social anxiety in middle childhood: link with behavioural inhibition when young
- 28 Review: Cannabis use increases the risk of psychotic outcomes
- 29 Maternal affective or substance disorders are risk factors for subsequent pregnancy loss
- 30 Reduced gestational age associated with an increased likelihood of depression in later life

Prevalence

- 31 An estimated 3.4% of adults have adult attention-deficit hyperactivity disorder
- 32 Autism and prenatal exposure to storms

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

 Articles carrying the Unlocked Logo are freely available online under the BMJ Journals unlocked scheme. See <http://ebmh.bmj.com/info/unlocked.dtl>

 www.publicationethics.org.uk
This journal is a member of and subscribes to the principles of the Committee on Publication Ethics