

Evidence-Based Mental Health



Contents

May 2008 Vol 11 No 2

EBMH Notebook

- 33** Nothing new under the sun
S Reid
- 34** Antidepressants and suicide symptoms: compelling new insights from the FDA's analysis of individual patient level data
C Barbui, A Cipriani, J R Geddes
- 36** Appraising the evidence: what is measurement bias?
L A Page, M Henderson
- 38** Et al

Prognosis

- 40** Problems identifying odours predicts mild cognitive impairment in the elderly
- 41** Attention deficit hyperactivity disorder is associated with poorer academic performance

Therapeutics

- 42** Cognitive improvements with antipsychotics: real or practice effect?
- 43** Ongoing pharmacotherapy maintains quality of life benefits in older adults with major depression
- 44** Depression in older adults: collaborative care model seems effective
- 45** Review: Intensive case management for severe mental illness reduces rehospitalisation when previous hospital use has been high
- 46** Depressed inpatients: treatment with psychotherapy plus pharmacotherapy superior to standard treatment
- 47** Telephone treatment support improves outcomes for depressed employees
- 48** Citalopram-resistant depression: cognitive therapy and medication similarly effective as second line treatments
- 49** Review: Psychotherapy for adolescents with depression: initial but no sustainable benefits
- 50** Antisocial behaviour: does a preventative intervention reduce cortisol levels in response to a social challenge?

- 51** Personalised feedback on alcohol problems in young adults is more effective as part of a motivational interview
- 52** Individual placement and support programme increases rates of obtaining employment in people with severe mental illness
- 53** Review: Psychological interventions for caregivers of people with dementia

Aetiology

- 54** New use of antipsychotic drugs in elderly people with dementia may increase the mortality risk
- 55** Offspring of suicide attempters at greater risk of suicide events
- 56** More severe symptoms of depression increase coronary heart disease mortality

Prevalence


- 57** Depression: greater effect on overall health than angina, arthritis, asthma or diabetes
- 58** Major depressive disorder is common in the elderly population of Yoruba-speaking regions of Nigeria
- 59** Major depressive disorder: afternoon and evening diurnal mood variation is common
- 60** Increase in out-based physician diagnoses of bipolar disorder in youth in the USA
- 61** Review: Black and minority ethnic people are more likely to be detained under the Mental Health Act 1983—no clear evidence why
- 62** Metabolic syndrome common in people with psychotic disorders, particularly in users of high potency antipsychotics
- 63** Incidence of schizophrenia lower in people with type I diabetes

Miscellaneous

- 64** Glossary

Electronic pages

- e1** Purpose and procedure

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.



www.publicationethics.org.uk
This journal is a member of and subscribes to the principles of the Committee on Publication Ethics