


Evidence-Based Mental Health



Contents

August 2008 Vol 11 No 3

EBMH Notebook

- 65 ebmh-talk *S Reid*
- 65 So careless of the single trial *P Tyrer*
- 66 Do antidepressants work? A commentary on "Initial severity and antidepressant benefits: a meta-analysis of data submitted to the Food and Drug Administration" by Kirsch et al
R H McAllister-Williams
- 68 A letter from America: rescuing inpatient psychiatry *J C Markowitz*
- 69 *Et al*

Prognosis

- 72 Prognostic factors for progression to psychosis in high risk youth
- 73 Mortality in people with dementia taking antipsychotics higher than in those taking other psychiatric medications

Therapeutics

- 74 Collaborative consultation increases use of medication titration trials but not behaviour in children with ADHD
- 75 No long-lasting effects of classroom-based cognitive behavioural and interpersonal prevention programmes on depressive symptoms in adolescents
- 76 Review: Cognitive behavioural therapy for adolescents with depression
- 77 Treatment-resistant depression: adding low dose risperidone improves symptoms
- 78 Adding repetitive transcranial magnetic stimulation to antidepressants does not improve response in people with depression
- 79 Review: Psychosocial and psychological interventions reduce postpartum depressive symptoms
- 80 Six-month multicomponent intervention improves postnatal depression in low-income settings
- 81 Review: Children and adolescents with schizophrenia spectrum disorders respond to antipsychotics, but are susceptible to adverse events
- 82 Metformin and lifestyle changes reduce antipsychotic-related weight gain in adults with schizophrenia

- 83 Occupational therapy is cost-effective for older people with dementia and their caregivers
- 84 Donepezil is no more effective than placebo for agitation in people with Alzheimer's disease
- 85 Antipsychotics for aggressive behaviour: no better than placebo for non-psychotic adults with learning difficulties
- 86 Intramuscular haloperidol plus promethazine is more effective and safer than haloperidol alone for rapid tranquillisation of agitated mentally ill patients
- 87 Tranquillising agitated mentally ill patients: intramuscular haloperidol plus promethazine reduces need for additional medication or medical attention compared to olanzapine
- 88 Review: Screening and intervention for alcohol misuse in emergency rooms reduces alcohol-related injuries
- 89 Structured patient-clinician communication using DIALOG improves patient quality of life
- 90 Review: Psychotherapy may be beneficial for people with hypochondriasis

Aetiology

- 91 Post-traumatic stress disorder increases the risk of new drug problems in young adults
- 92 Sudden infant death syndrome is associated with parental psychiatric admission
- 93 Schizophrenia associated with increased risk of colon cancer but reduced risk of respiratory cancer

Prevalence

- 94 Immigrants in areas of low immigrant density have a greater incidence of psychotic disorders than native Dutch
- 95 Increased risk of sexual offending in men with psychotic disorders

Miscellaneous

- 96 Glossary

Electronic pages

- e1 Purpose and procedure

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics
www.publicationethics.org.uk

 equator
network