

Evidence-Based Mental Health



Contents

February 2010 Vol 13 No 1


EBMH Notebook

- 1 Treating eating disorders: a review of the evidence
Rano Bhadoria, Kate Webb, John F Morgan
- 5 Maximising the treatment outcomes of opioid substitution treatment
Ed Day
- 8 Et al


Prognosis

- 9 Attention problems in primary school affect education achievements at high school

Diagnosis


- 10 Conners' Teacher Rating Scale has limited ability to predict DSM-IV ADHD in referred schoolchildren
- 11  New Alzheimer's disease screening test performs well in UK outpatient settings

Therapeutics

- 12 Video decision support affects patients' care choices in dementia care planning
- 13 Antidepressants combined with self-management of pain improves outcomes in people with comorbid pain and depression
- 14  Health visitor training reduces risk of postnatal depression 6 months after birth
- 15 School based prevention programme may reduce depressive episodes in adolescents at risk
- 16 Group CBT prevents depression in adolescents
- 17 Group therapy does not reduce repeated deliberate self-harm in adolescents
- 18 Group CBT reduces child anxiety diagnoses compared with non-specific group support


- 19 Review: pharmacotherapy increases response and reduces symptom severity in paediatric anxiety disorders
- 20 CBT improves worry severity in older adults with GAD in primary care
- 21 Neurofeedback training improves ADHD symptoms more than attention skills training
- 22 Citalopram not effective for repetitive behaviour in autistic spectrum disorders
- 23 Haloperidol plus promethazine effective for psychosis induced aggression
- 24 Second generation antipsychotics reduce dropout rates compared with first generation antipsychotics
- 25 Subcutaneous naltrexone implants reduce opioid use in opiate dependent patients
- 26 Review: methadone increases retention and reduces heroin use compared with non-pharmacological maintenance
- 27 Review: buprenorphine better than α_2 adrenergic agonists for managing opioid withdrawal
- 28 Adding zolpidem to CBT produces limited benefits in persistent insomnia
- 29 Review: multiple session early psychological interventions after trauma do not prevent PTSD

Aetiology

- 30 Long term antidepressant use linked to increased risk of diabetes
- 31  Schizophrenia and comorbid substance abuse substantially increases risks of violent crime

Prevalence

- 32 Estimated prevalence of autism spectrum conditions in Cambridgeshire is over 1%

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org

 equator network

 recycle
When you have finished with this please recycle it