


Evidence-Based Mental Health



Contents

May 2010 Vol 13 No 2

Purpose and procedure

33

EBMH Notebook

35 Qualitative Research Methods in Mental Health



Dr Sarah Peters

Prognosis

- 41 Review: depression is associated with increased cancer mortality
- 42 Girls with ADHD have an increased risk of psychopathology in young adulthood
- 43 Recurrence of bipolar spectrum disorders—a common issue among the affected population of young people
- 44 Anticholinergic drugs increase the risk of cognitive decline and dementia in older people
- 45 Depressive symptoms highly prevalent after a musculoskeletal workplace injury
- 46 No differences in mortality and suicide between treated and never-treated people with schizophrenia
- 47 Failure to improve during first 6 months of treatment and depression severity predict increased mortality in people with ACS and MDD

Therapeutics

- 48 Interpersonal therapy or guided self-help CBT improve remission from binge eating compared to a behavioural weight-loss programme at 2-year post-treatment
- 49 Review: benefits of antidepressants over placebo limited except in very severe depression
- 50 Online cognitive behavioural therapy is effective for depression in primary care
- 51 Review: Is aripiprazole as effective as risperidone in people with schizophrenia?
- 52 Review: antidepressants associated with increased risk of suicidality in adults aged less than 25 years

- 53 Review: ziprasidone is marginally less effective than other atypical antipsychotics in people with schizophrenia
- 54 Review: most antipsychotic drugs more than double the prolactin levels in children and adolescents
- 55 Review: There is no evidence that psychosocial interventions prevent disability after traumatic physical injury
- 56 Review: lack of evidence for the optimal dose of risperidone for schizophrenia

Aetiology

- 57 Estimates of how many cannabis users need to be prevented in order to prevent one case of schizophrenia
- 58 Increased risk of septal heart defects in newborns as a result of sertraline and citalopram intake during pregnancy
- 59 Frequent childhood victimisation predicts later psychiatric problems in females
- 60 Antipsychotic treatment associated with higher risk of hyperglycaemia in older persons with diabetes

Prevalence

- 61 Increase in proportion of patients on long-term medication accounts for increase in antidepressant prescribing in UK
- 62 Increased antidepressant and antipsychotic use in the USA between 1996 and 2005
- 63 HIV-positive young people at high risk of psychiatric disorders

Treatment

- 64 Limited evidence that antipsychotic drug treatment is associated with reduced brain volume

Glossary

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org

 equator network

 recycle
When you have finished with this please recycle it