


Evidence-Based Mental Health



Contents

May 2011 Vol 14 No 2

Purpose and procedure

- 33 Purpose and procedure

EBMH Notebook

- 35 Antipsychotics in dementia – mortality risks and strategies to reduce prescribing

Aetiology

- 37 Greater mortality risk with comorbid coronary heart disease and depressive symptoms either condition alone
- 38 Likelihood of later hospital admission for deliberate self-harm may be influenced by various risk factors in young people smoking
- 39 Patients receiving opiate substitution have elevated mortality risks at the start and end of treatment
- 40 Review: adverse effects of antidepressants use during pregnancy
- 41 Lower plasma β -amyloid levels are associated with moderately greater rate of cognitive decline among older people without dementia

Prevalence

- 42 The 2004 FDA warning on antidepressant suicidality risk changed outpatient treatment patterns for young people with new-onset depression
- 43 Increased adolescent mortality in the year after a psychiatric admission
- 44 Several medical and psychiatric conditions are more common in children and adolescents with bipolar disorder than the general population

Prognosis

- 45 Bereavement an independent risk factor for psychiatric symptoms in young people

- 46 Relationship between type 2 diabetes and risk of depression is bi-directional

Therapeutics

- 47 Review: metabolic side effects of second-generation antipsychotics
- 48 Review: reboxetine is ineffective and potentially harmful for acute treatment of major depression
- 49 A brief parent-focused intervention reduces anxiety disorders in socially inhibited children
- 50 Review: early intervention services can be clinically beneficial for people with early psychosis
- 51 Psychosis and comorbid substance misuse: integrated motivational interviewing and cognitive behavioural therapy reduces alcohol intake
- 52 Cognitive training for older drivers can reduce the frequency of involvement in motor vehicle collisions
- 53 Review: clozapine shows some improved clinical efficacy over risperidone and zotepine in treatment of schizophrenia, but robust evidence is lacking
- 54 For toddlers with autism spectrum disorders, supplementing a comprehensive intervention with interpersonal synchrony improves socially engaged imitation
- 55 Review: metformin is effective at attenuating the metabolic risks associated with use of atypical antipsychotics
- 56 Trauma-focused CBT reduces anxiety and post-traumatic stress disorder in children exposed to intimate partner violence

Glossary

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org

 equator network

 recycle
When you have finished with this please recycle it