


Evidence-Based Mental Health



Contents

February 2012 Vol 15 No 1

Editorial

- 1 Editorial

EBMH Notebook

- 2 Agomelatine and the brave old world of narrative-based medicine

Letter

- 4 Comment on risk assessment: predicting violence

Response to letter

- 5 Predicting violence – a response to Large and Olav

Aetiology

- 6 Review: depression is associated with an increased risk of stroke in adults
- 7 Antenatal depression associated with increased risk of offspring maltreatment and psychopathology

Prevalence

- 8 In England, a peak in poor mental health in middle-aged people is present only among low-income households
- 9 High rates of depressive symptomatology among injecting drug users in Saskatoon, Canada
- 10 Adult-onset chronic physical conditions are more common in people who report childhood adversities and early onset mental disorders
- 11 2.64% of South Korean children aged 7 to 12 have autism spectrum disorders
- 12 Review: some evidence of impaired neurocognitive performance in children and adolescents with bipolar disorder

Prognosis

- 13 Presence of psychiatric disorders is associated with poorer colon cancer outcomes in older adults
- 14 Mortality gap between people with schizophrenia or bipolar disorder and the general population persists in England
- 15 Only 10% of people with schizophrenia show sustained improvements over 3 years

Therapeutics


- 16 Review: some universal family-based prevention programmes provide small reductions in alcohol use in youths in the short and long term
- 17 Sertraline and mirtazapine do not reduce severity of depression in people with dementia
- 18 Trauma focused CBT for PTSD in young children is feasible, and may reduce PTSD symptoms
- 19 Review: second-generation antipsychotics improve response in paediatric bipolar disorder, but are associated with adverse events
- 20 Review: psychological and pharmacological treatments reduce depression symptoms in adults with coronary artery disease, but have no effect on mortality or cardiac events
- 21 Review: supportive interventions may improve short-term psychological distress in informal caregivers of patients at the end of life
- 22 Adjunctive cognitive behavioural therapy improves response in children with OCD and partial response to serotonin reuptake inhibitors

Treatment

- 23 Review: more research needed on early intervention for psychosis

Electronic page

- e2 Purpose and procedure

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org

 equator network

 recycle
When you have finished with this please recycle it