


# Evidence-Based Mental Health



## Contents

May 2012 Vol 15 No 2

### Editorial

- 25 Evidence-based mental health and psychosocial support in humanitarian settings: gaps and opportunities

### Letter

- 27 Response to Dr Goy's commentary

### Aetiology

- 28 Childhood adversity and certain mental health disorders are associated with increased risk of incident drug use among adults
- 29 Childhood and young adult-onset depression are associated with similar psychosocial risk factors
- 30 Comorbid dyslipidaemia and diabetes, but not antipsychotic use, is associated with unexplained sudden death among psychiatric patients
- 31 Sibling recurrence rate of autism spectrum disorders is 18.7%
- 32 Review: depression is associated with an increased risk of developing stroke
- 33 Review: premorbid IQ is inversely associated with future risk of schizophrenia

### Diagnosis

- 34 The Strengths and Difficulties Questionnaire hyperactivity-inattention subscale is more sensitive for the ADHD-combined subtype than other subtypes in 7–9-year-old school children

### Prevalence

- 35 Poor psychological health and stressful-life events are more common in adolescents who self-harm
- 36 ADHD is associated with an increased risk of written-language disorder
- 37 Children with social phobia have distinct social skills deficit and vocal characteristics to children with Asperger's disorder and typically developing children

### Prognosis

- 38 Suicide and attempted suicide are more common in children and adolescents in care, but rates of attempted suicide are higher before entry into care than after

- 39 Depression 3 and 9 months after discharge is less common in cardiac patients who are receiving statins at discharge
- 40 Parental ADHD affects parent and child outcomes of parental friendship coaching
- 41 Regular primary care and specialty care as needed is associated with remission from alcohol and drug use disorders over 9 years
- 42 In Ethiopia, people with schizophrenia are at higher risk of 5-year mortality than the general population
- 43 Use of ADHD drugs in children and young adults does not increase risk of serious cardiovascular adverse events
- 44 The risk of suicide attempt by a child or adolescent is highest after a contact with a psychiatric department

### Therapeutics

- 45 Review: risperidone, olanzapine and haloperidol are the most effective drugs for acute mania in adults with bipolar I disorder
- 46 Review:  $\omega$ -3 fatty acids produce a small improvement in ADHD symptoms in children compared with placebo
- 47 Review: extrapyramidal side effects vary between different second-generation antipsychotic drugs
- 48 Multisystemic therapy reduces re-offending in young offenders between 12 and 18 months post-treatment
- 49 Online cognitive-behaviour therapy is similarly effective to clinic-based CBT for reducing adolescent anxiety
- 50 Increasing psychotherapy dose from 8 to 16 sessions does not improve social functioning in people with major depressive disorder receiving pharmacotherapy
- 51 Review: limited evidence from two small trials suggests no improvement in ASD symptoms with short term  $\omega$ -3 supplementation
- 52 The Strongest Families intervention is more effective than usual care in children with mild or moderate oppositional, attention or anxiety disorders

### Electronic page

- e3 Purpose and procedure

This article has been chosen by the Editor to be of special interest or importance and is freely available online.

COPE COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

[www.publicationethics.org](http://www.publicationethics.org)

equator network

recycle

When you have finished with this please recycle it