


# Evidence-Based Mental Health



## Contents

November 2012 Vol 15 No 4

### Editorial

- 85 Policing and mental health

### Prognosis

- 86 Changes in occipital lobe gyrification may be an early marker for psychosis in youth with velo-cardiofacial Syndrome
- 87 Reported loneliness rather than social isolation is a risk factor for 10-year mortality in older men

### Therapeutics

- 88 Modular psychotherapy improves problems in youth with anxiety, depression or conduct disorder more rapidly than standard psychotherapy
- 89 Cognitive behavioural treatments have long-term benefits in female rape survivors with post-traumatic stress disorder
- 90 SPARX computerised CBT is as effective as usual care for mild-to-moderate depression in help seeking adolescents
- 91 No difference in extrapyramidal side effects between first-generation and second-generation antipsychotics
- 92 Review: maintenance antipsychotics prevent relapse in patients with schizophrenia
- 93 Adding facilitated physical activity to standard GP care for depression does not improve symptoms
- 94 Review: urgent need for RCT evidence on effectiveness of crisis interventions for borderline personality disorder
- 95 Community-based outreach may reduce distress in women exposed to intimate partner abuse more than system-based referral
- 96 Atomoxetine reduces ADHD symptoms in children with autism spectrum disorder

### Aetiology


- 97 Use of benzodiazepines in adults with schizophrenia is associated with increased mortality
- 98 Fluoxetine and sertraline may be associated with lower risk of suicide death than paroxetine in adults with depression
- 99 Veterans who have served in the conflicts of Iraq and Afghanistan are at increased risk of suicide compared to other veterans if they have a psychiatric diagnosis
- 100 Review: adverse effects of different second-generation antipsychotics vary in children and adolescents
- 101 Following self-harm, there are shared and differing risk factors for subsequent suicide death or accidental death
- 102 Hyperactivity/impulsivity in childhood, but not inattention, is associated with early-onset tobacco and alcohol use

### Prevalence

- 103 Review: adults with disabilities experience more violence than non-disabled adults
- 104 Higher BMI, lower self-esteem and more abnormal eating habits are associated with greater body dissatisfaction in adolescents
- 105 Prevalence of psychotic symptoms in the general population varies across 52 countries
- 106 Review: emotion identification deficits are associated with functional impairments in people with schizophrenia
- 108 About 7% of 4-year-olds in Norway meet criteria for a psychiatric diagnosis

### Electronic page

- e4 Purpose and procedure

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

**C O P E** COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

[www.publicationethics.org](http://www.publicationethics.org)

 equator network

 recycle  
When you have finished with this please recycle it