


Evidence-Based Mental Health



Contents

February 2013 Vol 16 No 1

Editorial

- 1 Student mental health; below the radar?



EDITOR'S CHOICE

Expert review

- 2 Drop the language of disorder



EDITOR'S CHOICE

Prognosis

- 4 Low-level alcohol consumption in early pregnancy may not affect child intelligence, attention or executive function at 5 years of age
- 5 Boys with ADHD are at increased risk of psychosocial, educational and functional impairment in adulthood

Therapeutics

- 6 A primary care-based treatment programme improves postpartum depression at 12 months
- 7 Tuberculosis patients in South Africa have a high prevalence of psychological distress
- 8 Targeted education for general practitioners reduces risk of depression or suicide ideation or attempts in older primary care patients
- 9 A preventative intervention for disruptive young boys may reduce adult criminality by reducing antisocial behaviour in adolescence
- 10 Review: non-steroidal anti-inflammatory drugs may reduce schizophrenia symptom severity in the short term when added to antipsychotics
- 11 Review: social skills groups may improve social competence in children and adolescents with autism spectrum disorder
- 12 Review: little evidence that PUFA supplementation improves symptoms in ADHD
- 13 The Family-School Success intervention improves some family and educational outcomes in children with attention deficit hyperactivity disorder more than a control psychosocial intervention
- 14 Review: limited evidence suggests mental health literacy interventions may improve help-seeking attitudes for depression, anxiety and general psychological distress
- 15 The DAISY psychosocial intervention does not improve outcomes in patients with mild Alzheimer's disease or their carers

- 16 Similar benefits with dialectical behaviour therapy or general psychiatric management for people with borderline personality disorder
- 17 Review: some evidence of benefit for psychotherapies in borderline personality disorder
- 18 Non-pharmacological interventions reduce antipsychotic-associated weight gain in outpatients
- 19 Long-acting risperidone increases medication costs and does not improve outcomes in unstable schizophrenia
- 20 An assertive outreach intervention does not reduce repeat suicide attempts compared with usual care
- 21 Review: antipsychotics do not increase weight in women with anorexia nervosa
- 22 Review: non-pharmacological interventions delivered by family caregivers improve symptoms in people with dementia

Aetiology

- 23 Perceived daily psychological demands in healthy adults not taking blood pressure medication are associated with progression of carotid atherosclerosis
- 24 Children of migrant parents may be at greater risk of low-functioning autism spectrum disorder, but lower risk of high-functioning autism spectrum disorder
- 25 Boys born small for gestational age may be at an increased risk of symptoms of depression and anxiety during adolescence if exposed to chronic stress
- 26 Higher dietary intakes of potassium, calcium and magnesium are associated with a reduced risk of developing vascular dementia
- 27 Past exposure to neuroleptics is associated with the risk of developing Parkinson's disease

Prevalence

- 28 Self-harm is common in adolescents in England

Receive regular table of contents by email.
Register using this QR code.



This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org

equator
network

recycle

When you have finished with this please recycle it