


# Evidence-Based Mental Health



## Contents

November 2014 Vol 17 No 4

### Clinical review

- 97 Self-harm in young people



### Causes and risk factors

- 100 Young maternal age and old paternal age induce similar risk of mental disorders in offspring
- 101 High rates of suicide attempt in early-onset psychosis are associated with depression, anxiety and previous self-harm
- 102 Medication reduces crash risk among men with ADHD

### Evidence-based case conference

- 103 Adverse effects of antidepressants during pregnancy



### Prevalence, assessment and diagnosis

- 105 Developing accuracy when estimating Global Burden of Disease (GBD) in mental and substance use disorders using complex research methodology, refined data analysis and community participation
- 106 Neonatal factors associated with autism spectrum disorders in infants
- 107 Children exposed to trauma should be screened for symptoms of PTSD

### Pharmacological interventions

- 108 Atomoxetine is a second-line medication treatment option for ADHD
- 109 Ziprasidone monotherapy is ineffective for bipolar-associated anxiety
- 110 Paliperidone depot is not different from haloperidol in relapse prevention of schizophrenia, but different side effects should be considered

### Statistics in practice

- 111 A hands-on practical tutorial on performing meta-analysis with Stata




### Psychological interventions

- 117 Extending the evidence base for prolonged exposure: significant effectiveness noted with sexual abuse-related PTSD in adolescent girls
- 118 Finally moving beyond the horse race: CBT and psychodynamic therapy equally effective for depression
- 119 Some evidence of efficacy but CBT not the answer to mental health needs in general hospitals

### Miscellaneous

- 120 Theory of mind interventions can be effective in treating autism, although long-term success remains unproven
- 121 Targeted, preventive interventions in primary school prevent mental health problems in adolescence
- 122 Sleep therapy is effective in reducing symptoms of depression among adolescents
- 123 Review tempers expectations for peer-led services but further studies are required
- 124 Community-based intervention for people with schizophrenia successfully provided by supervised community health workers in a low-resource setting

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

[www.publicationethics.org](http://www.publicationethics.org)

 equator network

 recycle  
When you have finished with this please recycle it

Receive regular table of contents by email.  
Register using this QR code.

