


# Evidence-Based Mental Health



## Contents

February 2015 Vol 18 No 1

### Clinical review

- 1 Maintaining mood stability in bipolar disorder: a clinical perspective on pharmacotherapy



### Perspective

- 7 Early identification of recurrent mood disorders in youth: the importance of a developmental approach



### Causes and risk factors

- 10 Mixed associations observed between advanced paternal age and psychiatric morbidity in offspring
- 11 Early antecedents of suicide: the role of prenatal and childhood risk factors

### Prevalence, assessment and diagnosis

- 12 One-third of Danish people will receive secondary care treatment for a mental disorder during their lifetime
- 13 Identifying inmates that will actually die by suicide

### Evidence-based case conference

- 14 Helping people with schizophrenia to quit smoking

### Pharmacological interventions

- 16 Study presents limited assessment of pharmacotherapy for alcohol use disorders
- 17 No better solution for antipsychotic treatment with regard to long-term safety and effectiveness

### Psychological interventions

- 18 Psychosis patients refusing antipsychotic medicine could benefit from CBT in terms of both symptom reduction and social functioning

- 19 Group-based parent training programmes demonstrate positive effects on the psychosocial well-being of both mothers and fathers
- 20 Day hospital treatment following brief inpatient medical stabilisation for adolescents with anorexia nervosa is equivalent in efficacy to inpatient treatment
- 21 Seven-day intensive cognitive therapy for PTSD is as effective as weekly cognitive therapy and more effective than weekly supportive therapy
- 22 Developing personalised integrated psychotherapy for patients with personality disorders

### Statistics in practice

- 23 Accounting for baseline differences in meta-analysis



### Miscellaneous

- 27 Smartphone applications can help in treatment for alcoholism
- 28 Depressed with cancer can respond to antidepressants, but further research is needed to confirm and expand on these findings
- 29 Management of dementia-related agitation: between the devil and the deep blue sea
- 30 Time efficient interventions by general practitioners curb benzodiazepine consumption among long-term users
- 32 Cultural adaptations of psychological treatments for depression are mostly based on implementation rather than content

### Correction

- 26 Correction

This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics  
[www.publicationethics.org](http://www.publicationethics.org)

equator network

recycle  
When you have finished with this please recycle it

Receive regular table of contents by email.  
Register using this QR code.

