


Evidence-Based Mental Health



Contents

August 2015 Vol 18 No 3

Expert commentary

- 65** Intervening early in children with bipolar disorder: is there a pot at the end of the Rainbow?


Systematic review

- 67** Parenting in mothers with borderline personality disorder and impact on child outcomes


Forensic psychiatry

- 76** The Penrose hypothesis in the 21st century: revisiting the asylum


Clinical review

- 77** Increasing placebo response in antipsychotic trials: a clinical perspective

Evidence-based case conference

- 80** Which drug is the best bet when your bipolar patient gets depressed?

Causes and risk factors

- 82** SSRI antidepressants are not associated with cerebral microbleeds or ischaemic vascular lesions
- 83** Sibling bullying is associated with anxiety, depression and self-harm
- 84** Environmental factors, including cannabis, are strongly related to the age of onset and morbidity of schizophrenia
- 85** Comorbid tics have no effect on response to cognitive-behavioural therapy in youth with obsessive-compulsive disorder
- 86** Monitoring of suicide risk throughout the course of treatment with antidepressants for depression is required, but vigilance is required for those on some particular antidepressant agents

Outcomes

- 87** Understanding the course of cognitive deficits over the onset of psychosis

Pharmacological interventions

- 88** Pharmacotherapy for bipolar depression: comparative efficacy and acceptability is in the eye of the beholder
- 89** Metformin to reduce weight gain and metabolic disturbance in schizophrenia

Psychological interventions

- 90** Efficacy of complicated grief therapy in older adults
- 91** Psychosocial therapy after self-harm associated with reduced repetition, suicide, and all-cause mortality in Denmark
- 92** Computerised cognitive behavioural therapy has the potential to prevent and treat anxiety and depression in young people
- 93** Some promising news about psychosocial interventions for adolescent self-harm

Miscellaneous

- 94** Integrating depression treatment within primary care improves outcomes in adolescents
- 95** Cognitive therapy plus medication management is better than antidepressants alone for patients with severe depression
- 96** Web-based prevention and treatment programmes for internalising problems in youth: encouraging findings in an expanding research field

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org

 equator network

 recycle
When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

