


Evidence-Based Mental Health



Contents

February 2016 Vol 19 No 1

Perspectives

- 1** Is treatment-resistant depression a useful concept?
 EDITOR'S CHOICE
- 4** Control or overcontrol for covariates?

Clinical reviews

- 6** Vitamin D in schizophrenia: a clinical review
- 10** Predicting the onset of psychosis in patients at clinical high risk: practical guide to probabilistic prognostic reasoning
 EDITOR'S CHOICE

Statistics in practice

- 16** Approaches for synthesising complex mental health interventions in meta-analysis
 EDITOR'S CHOICE

Forensic psychiatry

- 22** Access to firearms in the USA: angry and impulsive behaviour in people with and without mental disorders

Causes and risk factors

- 23** Bullying victimisation increases the risk of emergence and persistence of specific psychotic symptoms and of psychotic disorder
- 24** Antidepressants are associated with hospital admitted intracranial bleeds in people taking other medication associated with bleeding

Pharmacological interventions

- 25** Examination of a four-step treatment algorithm for postpartum psychosis
- 26** SSRIs have a smaller benefit in paediatric when compared to adult major depressive disorder

Psychological interventions

- 27** Brief and effective psychological treatment of worry within people diagnosed with persecutory delusions
- 28** Targeting social and non-social cognition to improve cognitive remediation outcomes in schizophrenia
- 29** Cross-cultural efficacy of the Cool Kids programme for child and adolescent anxiety

Miscellaneous

- 30** Preschool hyperactivity increases costs into early adulthood
- 31** Health promotion coaching for obese individuals with serious mental illnesses produces clinically significant reductions in cardiovascular risks up to 18 months

Letter

- 32** Comment on: 'Reports of recovery in chronic fatigue syndrome may present less than meets the eye'

Electronic pages

- e1** Mediterranean diet and treating diabetes and depression in old age may reduce dementia risk
- e2** Cognitive-behavioural therapy for insomnia: effective, long-lasting and safe
- e3** Dropout in self-guided web-based interventions for depression can be predicted by several variables
- e4** Promoting multilevel primary prevention of depression and diabetes during midlife may protect against dementia
- e5** Antidepressant use with D-Cycloserine may block fear extinction

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

 This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebmh.bmj.com/site/about/guidelines.xhtml#open>

 This journal is a member of and subscribes to the principles of the Committee on Publication Ethics www.publicationethics.org

 equator network

 recycle
When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

