


Evidence-Based Mental Health



Contents

November 2016 Vol 19 No 4

Perspective

- 97 Methylphenidate for ADHD in children and adolescents: throwing the baby out with the bathwater
- 100 Evidence on methylphenidate in children and adolescents with ADHD is in fact of 'very low quality'

Clinical reviews

- 103 Exercising control over bipolar disorder
- 106 Assessing and measuring cognitive function in major depressive disorder

Old age psychiatry

- 110 Driving and dementia: a clinical update for mental health professionals

Statistics in practice

- 114 Designing and analysing clinical trials in mental health: an evidence synthesis approach

Evidence-based case conference

- 118 Which psychotherapy for PTSD?

Commentaries

Causes and risk factors

- 120 PTSD is associated with elevated inflammation: any impact on clinical practice?

Outcomes

- 121 Prevention and treatment of mental health crises: examining the gaps in the evidence base
- 122 Perinatal suicide associated with depression diagnosis and absence of active treatment in 15-year UK national inquiry

Pharmacological interventions

- 123 Routine use of antipsychotics to prevent or treat delirium is not recommended

- 124 Metoclopramide may be effective for clozapine-induced hypersalivation

Psychological interventions

- 125 No evidence that CBT is less effective than antidepressants in moderate to severe depression

Miscellaneous

- 126 Does community-based suicide prevention decrease suicide attempts? Strategies for reaching zero suicide
- 127 Second-generation antidepressants and cognitive-behavioural therapy are both viable choices for initial treatment of major depression

Letter

- 128 Are molecular scanners going to make double-blind placebo-controlled trials impossible?

Further commentaries

Electronic pages

- e22 Reconsidering risk factors for renal dysfunction in lithium-treated patients
- e23 Cognitive impairment in depression and its (non-)response to antidepressant treatment
- e24 Lack of clinically useful response predictors for treating aggression and agitation in Alzheimer's disease with citalopram
- e25 Is long-acting injectable aripiprazole useful for the treatment of acute exacerbation of schizophrenia?
- e26 Are sertraline, paroxetine and duloxetine the most effective antidepressants for use in depressed adults over 60 years?
- e27 Autoimmune diseases can be associated with depression
- e28 Reduction in general intelligence and executive function persists into adulthood among very preterm or very low birthweight children

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

 This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebmh.bmj.com/site/about/guidelines.xhtml#open>

 This journal is a member of and subscribes to the principles of the Committee on Publication Ethics www.publicationethics.org

 equator network

 recycle
When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

