

Evidence-Based Mental Health

Contents

Purpose and procedure	66	Review: haloperidol decanoate is superior to placebo for schizophrenia	84
EBMH notebook		Nortriptyline plus interpersonal psychotherapy was effective in major depression in older adults	85
Teaching evidence-based practice in mental health	68	Traditional child psychotherapy did not reduce psychopathology compared with academic tutoring ...	86
The effect of misclassification on estimates of relative risk	70	Bright light and high density negative air ionisation reduced symptoms in seasonal affective disorder	87
Other articles noted	72	Bright morning light reduced depressive symptoms in seasonal affective disorder	88
Therapeutics		Review: medication and cognitive behaviour therapy control symptoms of bulimia nervosa	89
A nicotine patch plus nicotine nasal spray was more effective than a nicotine patch alone for smoking cessation.....	73	Diagnosis	
Bupropion alone or with a nicotine patch increased smoking cessation rates	74	CAMCOG detected dementia and cognitive impairment in Parkinson's disease	90
Lamotrigine for bipolar I disorder improved depressive symptoms on some depression scales	75	Clinical criteria for three types of dementia had low sensitivity and high specificity	91
CBT plus parental anxiety management was more effective when the child and parent(s) were anxious	76	Prognosis	
Antihypertensive treatment may have reduced the rate of dementia in older patients with isolated systolic hypertension	77	Behavioural problems in infants and toddlers were associated with later psychiatric disorders	92
Review: antidepressants improve depression in adults with physical illnesses	78	Review: in non-referred samples of children, early externalising behaviour does not predict later antisocial behaviour	93
Befriending led to a higher frequency of overall remission in women with chronic depression	79	Aetiology	
Cognitive behavioural group therapy and phenelzine were both effective in social phobia	80	APOE ϵ 4 allele, age, and duration of unconsciousness were associated with unfavourable outcomes in traumatic brain injury	94
Review: psychosocial treatments can play an important part in the comprehensive management of patients with schizophrenia	81	Review: psychotropic drugs increase the risk of falls in older people	95
Review: <i>Ginkgo biloba</i> extract improves cognitive function in mild to moderate Alzheimer's disease	82	Family disadvantages in early life were associated with depression in adulthood	96
Teaching patients to recognise early symptoms of relapse reduced manic relapses and improved social functioning in bipolar disorder	83	Glossary	Inside back cover