

Evidence-Based Mental Health

Contents

Purpose and procedure (abbreviated)	98
EBMH notebook	
Evaluating qualitative research	99
Evidence-based medicine and complementary medicine	102
Other articles noted	104
Therapeutics	
A very brief 3 session psychotherapy intervention was effective in subsyndromal depression	105
Review: lower doses of antidepressant drugs are effective and have fewer adverse effects in depression ..	106
Intravenous lorazepam reduced alcohol related seizures in patients with chronic alcohol abuse	107
Equivalent response to methylphenidate in children with ADHD plus anxiety and those with ADHD only ..	108
Slow release morphine did not reduce neonatal abstinence syndrome more than methadone	109
Brief physician advice reduced drinking in older adults ..	110
Coping skills training was effective in adolescents with type 1 diabetes	111
Eptastigmine led to cognitive, clinical, and functional benefits in Alzheimer's disease	112
Memantine led to functional improvement and reduced care dependence in severe dementia	113
Slow, repetitive transcranial magnetic stimulation was effective for major depression	114
Review: greater intensity of light leads to a greater reduction in typical (but not atypical) depressive symptoms in seasonal affective disorder	115
Sertraline was effective in children and adolescents with obsessive compulsive disorder	116
Cognitive behaviour therapy or temazepam, or both, improved short term outcomes for older adults with persistent insomnia	117
Clozapine improved psychosis without worsening parkinsonism	118
Diagnosis	
NINCDS-ADRDA criteria were not able to exclude frontotemporal dementia	119
Prognosis	
1 year later, 9–13 year olds impaired but not diagnosed were as disturbed as those diagnosed.....	120
Performance on the praxis subscale of the CAMCOG was associated with survival in patients with Alzheimer's disease	121
Risk of homelessness was increased in schizophrenia when 3 factors were present	122
Most patients returned to previous occupational level 1 year after head injury	123
Aetiology	
A decrease in care was associated with suicide in mental illness	124
An inverted U relation existed between unemployment and foster home placements	125
Current cigarette smoking was associated with a higher risk of Alzheimer's disease	126
Older women with depression were at increased risk for falls and fractures	127
Economics	
Assertive community treatment for homeless adults with severe mental illnesses was effective but not more expensive than usual care	128
Subject index	129
Author index	134
Contents	
Glossary	Inside back cover