


Evidence-Based Mental Health



Contents

Perspective

- 1 Twittering on about mental health: is it worth the effort? *M Jayaram, L Moran, C Adams*

Clinical reviews

- 4 Pharmacological and non-pharmacological treatment of adults with ADHD: a meta-review *F De Crescenzo, S Cortese, N Adamo, L Janiri*
- 12 Current status of electroconvulsive therapy for mood disorders: a clinical review *D Kolar*

Old age psychiatry

- 15 Cognitive-behavioural therapy for non-motor symptoms of Parkinson's disease: a clinical review *I Koychev, D Okai*

Statistics in practice

- 21 Exploratory trials in mental health: anything to learn from other disciplines? *S Trelle*

Commentaries

Forensic psychiatry

- 25 Impact of gun law reforms on rates of homicide, suicide and mass shootings in Australia *P Brown*

Outcomes

- 26 Involvement of patients in planning their future treatment may reduce compulsory admissions to hospital *S Kisely*
- 27 Evidence of ocular side effects of SSRIs and new warnings *J Kirkham, D Seitz*

Pharmacological interventions

- 28 Antipsychotic adjunctive therapy to mood stabiliser should be continued for 6 months after remission of a manic episode *K Hatta*

February 2017 Volume 20 Issue 1

Psychological interventions

- 29 Cognitive-behavioural group therapy for youth with high-functioning autism spectrum disorders demonstrates modest effects on social responsiveness *E A Storch*
- 30 Significant methodological flaws limit conclusions drawn by authors of a recent PTSD mindfulness study *D J Lee, C W Hoge*


Letters


- 31 Letter in response to Drs Lee and Hoge's commentary *C R Erbes, P Thuras, K O Lim, M A Polusny*
- 32 Randomised or not? *D Lufi, J Parish-Plass*

Further commentaries

Electronic pages

- e1 Prevention of depression and anxiety: is the whole better than the sum of the parts? *S Merry, S Hetrick*
- e2 Disruptive behaviour may hinder the acquisition of daily living skills for youth with autism spectrum disorder and disruptive behaviour *E A Laugeson*
- e3 More ACTIONS needed to reach a consensus on adjunctive antidepressant therapy for negative symptoms of schizophrenia *S Akhondzadeh, E Moazen-Zadeh*
- e4 Short web-based guided self-help intervention prevents the onset of a major depressive disorder in adults with subthreshold depression *L Bolier, B Boon*
- e5 Mindfulness-based cognitive therapy delays depressive relapse across demographic subgroups *G Meadows, F Shawyer*
- e6 Reduced all-cause mortality with antipsychotics and antidepressants compared to increased all-cause mortality with benzodiazepines in patients with schizophrenia observed in naturalistic treatment settings *J M Rubio, C U Corell*

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

 This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebmh.bmj.com/site/about/guidelines.xhtml#open>

 This journal is a member of and subscribes to the principles of the Committee on Publication Ethics www.publicationethics.org

 equator network

 recycle
When you have finished with this please recycle it

Receive regular table of contents by email.
Register using this QR code.

