


# Evidence-Based Mental Health



## Contents

May 2017 Volume 20 Issue 2

### Perspective

- 33 What is the impact of a research publication?  
*S Fazel, A Wolf*



OPEN ACCESS

### Clinical review

- 35 Psychosocial interventions for self-harm, suicidal ideation and suicide attempt in children and young people: What? How? Who? and Where?  
*G Cox, S Hetrick*



EDITOR'S CHOICE

### Original article

- 41 Implementing tools to support evidence-based practice: a survey and brief intervention study of the National Elf Service across Oxford Health NHS Foundation Trust  
*L Z Atkinson, A Forrest, L Marriner, J Geddes, A Cipriani*



EDITOR'S CHOICE

### Statistics in practice

- 46 Reporting guidance considerations from a statistical perspective: overview of tools to enhance the rigour of reporting of randomised trials and systematic reviews  
*B Hutton, D Wolfe, D Moher, L Shamseer*



EDITOR'S CHOICE

### Evidence-based case conference

- 53 Prognosis of delirium  
*T A Furukawa*

### Expert commentary

- 55 Immigrant women and women with learning disabilities have complex mental health needs and service use in the perinatal period  
*K Ayre, H Khalifeh*

### Commentaries

#### Causes and risk factors

- 58 Daily use of high-potency cannabis is associated with an increased risk of admission and more intervention after first-episode psychosis  
*M Large, O Nielssen*

#### Outcomes

- 59 Hypersomnia: an overlooked, but not overestimated, sleep disturbance in bipolar disorder  
*K A Kaplan, R Williams*

- 60 Relative to SSRI users, SSRI–statin users have fewer psychiatric hospital contacts and no increase in suicidal behaviour or all-cause mortality  
*C Andrade*

### Pharmacological interventions

- 61 Short-term adjunct of topiramate to antipsychotics in schizophrenia improves the psychopathology and has weight maintenance  
*T Kishi*

### Psychological interventions

- 62 Targeted psychological interventions may prevent depression in children and adolescents  
*E Watkins*
- 63 Simpler therapy may successfully treat adolescents with anorexia nervosa  
*S Agras*

### Letter

- 64 Changing response rates in clinical trials of depression: how did the introduction of DSM-III and DSM-III-R influence the outcome?  
*K N Fountoulakis*

### Further commentaries

#### Electronic pages

- e7 Largest study to date shows overall use of antipsychotics in pregnancy does not appear to significantly increase the risk of congenital malformations  
*S Grigoriadis, M Peer*
- e8 Uncertain association between depression and stroke risk in a Chinese mega-study  
*E J Brunner, I R Weinreb*
- e9 Mood Zoom could be a promising tool for daily mood variability monitoring, potentially differentiating bipolar from borderline patients  
*E Isometsä*
- e10 Cognitive–behavioural therapy can prevent transition to psychosis in ultra-high-risk participants in the long term  
*J-P Miron, A Abdel-Baki*

### Correction

- 54 Correction

This article has been chosen by the Editor to be of special interest or importance and is freely available online.

This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebmh.bmj.com/site/about/guidelines.xhtml#open>

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics [www.publicationethics.org](http://www.publicationethics.org)

equator network

recycle  
When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

