

# Evidence-Based Mental Health

## Contents

<b>Purpose and procedure</b> .....	2	Oral nalmefene reduced relapse to heavy drinking and was well tolerated in patients with alcohol dependence . . .	20
<b>EBMH notebook</b>		A behavioural intervention reduced depression and organic symptoms in elderly people in residential homes.....	21
Assessing allocation concealment and blinding in randomised controlled trials: why bother? .....	4	Review: cognitive and behaviour therapies are effective for chronic pain .....	22
Answering mental health questions with reliable research evidence .....	6	More people with depression continued treatment with fluoxetine than with desipramine or imipramine .....	23
<b>Other articles noted</b> .....	8	Review: nursing interventions increase smoking cessation rates in adults .....	24
<b>Therapeutics</b>		<b>Prognosis</b>	
A needs based intervention for patients with schizophrenia and their carers reduced relapse and carer needs .....	9	Failure to thrive in infancy was associated with later stunting and wasting but not cognitive or educational disadvantages .....	25
Review: rivastigmine may improve cognitive outcomes in Alzheimer's disease .....	10	Nature and outcome of psychotic illness was similar in all ethnic groups.....	26
Intensive case management was as effective as standard case management in severe psychotic illness .....	11	Pattern of course of psychosis and diagnostic stability did not differ by ethnic family background .....	27
Reframing memory reduced some measures of children's anticipatory distress and pain during lumbar puncture .....	12	Depressive coping and emotional distress predicted shortened survival in patients with lung cancer .....	28
Individual drug counselling plus group drug counselling was effective in patients with cocaine dependence.....	13	Patients with psychiatric disorders, particularly affective disorders, had an increased risk of developing dementia . . .	29
Individual placement and support led to more competitive employment than did enhanced vocational rehabilitation for severe mental disorder .....	14	<b>Aetiology</b>	
Review: studies with sufficient follow up do not show a clear benefit for pharmacotherapy in alcohol dependence.....	15	Review: schizophrenia is associated with premature rupture of membranes, gestational age < 37 weeks, and use of an incubator or resuscitation.....	30
Mentally ill adults had fewer homeless days in staffed group homes than in independent flats.....	16	Typical clinical doses of methylphenidate did not cause tics or exacerbate pre-existing tics in children with attention deficit hyperactivity disorder .....	31
A literacy promotion intervention increased the frequency of parent to child reading .....	17	<b>Qualitative</b>	
Cognitive behavioural therapy (CBT) with exercise led to higher levels of continuous smoking cessation in women than did CBT without exercise .....	18	Women with borderline personality disorder expressed they were living with a pejorative label, with self destructive behaviour viewed as manipulative, and with limited access to care.....	32
Review: progestogens do not resolve postnatal depression but oestrogens may improve depression scores.....	19	<b>Glossary</b> .....	Inside back cover