

Evidence-Based Mental Health

Contents

Purpose and procedure	34	Involuntary outpatient commitment did not reduce hospital recidivism, though may for some patient subgroups	53
EBMH notebook		Integrated outpatient treatment increased abstinence in men with alcohol related illness who were ongoing drinkers	54
Evidence based psychotherapy: special case or special pleading?	35		
Welcome to our new associate editors	37		
A new addition to this journal	37		
Resource corner	38		
Other articles noted	40		
Therapeutics			
Paroxetine was effective for reducing symptoms in social phobia	41		
The Fast Track prevention programme improved some outcomes in high risk schoolchildren	42		
The Fast Track prevention programme improved classroom atmosphere and peer reported behaviour problems	43		
Review: lithium augmentation increases treatment response in refractory depression	44		
Sertraline was more effective than desipramine or placebo for premenstrual syndrome	45		
Reading and social skills programmes had limited effectiveness on behavioural maladjustment in primary school children	46		
A community reinforcement and family training approach was effective in engaging unmotivated problem drinkers in treatment	47		
Antidepressants plus cognitive therapy reduced relapse in residual depression	48		
Plasma exchange and intravenous immunoglobulin improved symptoms in a subtype of obsessive compulsive disorder and tic disorders	49		
Review: psychotherapy may be of benefit in personality disorders	50		
Venlafaxine had higher response and remission rates than paroxetine in non-chronic treatment resistant depression	51		
Synthetic human secretin did not improve symptoms or communication skills in children with autistic disorders	52		
		Diagnosis	
		The 15 item Geriatric Depression Scale had perfect sensitivity but low specificity for detecting depression in elderly people	55
		The Even Briefer Assessment Scale for Depression in German was accurate for identifying depression in elderly patients	56
		Prognosis	
		Dementia in advanced age led to higher mortality rates and shortened life	57
		Aetiology	
		Review: most antipsychotic drugs are associated with weight gain	58
		A lower minimum legal drinking age was associated with increased suicide in youths 18–23 years of age.	59
		The association between cognitive function and mortality pertained to specific but not general measures of cognitive function when health factors were considered	60
		Review: 7.7 to 12.9 units of alcohol per week is the level of alcohol consumption at which mortality is the lowest in men	61
		Qualitative	
		Resilience was not necessarily a healthy concept for adolescents in inner city vocational high schools.	62
		Physicians indicated the need to frame questions and develop indirect approaches that foster patient trust in evaluating victims of domestic violence	63
		St John's Wort users reported: a history of exposure to herbal remedies and belief in their safety; depressed mood; and ease of access to alternative medicines	64
		Glossary	Inside back cover