

Evidence-Based Mental Health

Contents

Purpose and procedure	2	Behavioural family therapy reduced disruptive behaviour in children at risk for conduct problems	20
EBMH notebook		Review: nicotine replacement treatments achieve smoking abstinence at 6–12 months	21
So much research evidence, so little dissemination and uptake: mixing the useful with the pleasing	3	Cognitive behavioural therapy reduced distress and doctor visits in patients with medically unexplained symptoms	22
Applying the results of trials and systematic reviews to our individual patients	6	Review: newer drugs have not yet been shown to be better than clozapine for schizophrenia	23
Other articles noted	8	Review: donepezil, metrifonate, rivastigmine, and <i>Ginkgo biloba</i> are more effective than placebo in Alzheimer's disease	24
Therapeutics		Diagnosis	
Divalproex did not differ from placebo or lithium in preventing recurrence of new mood episodes in bipolar I disorder, remission phase	9	Review: strong evidence recommends genetic and metabolic testing in subgroups of children with autism	25
Review: family intervention added to standard treatment for schizophrenia reduces relapse	10	The Quick PsychoDiagnostics Panel was accurate for identifying psychiatric disorders in primary care	26
Review: patient education is effective for improving patient knowledge of schizophrenia	11	Prognosis	
Review: school based child sexual abuse prevention programmes are effective for improving prevention skills and knowledge	12	Review: the location of stroke is not associated with the risk of depression	27
Review: home visiting programmes that include ≥ 1 postnatal home visit are associated with improved quality of home environment and parenting	13	Diagnostic stability was higher in schizophrenia than other psychoses	28
A comprehensive coping strategy programme reduced nausea and fatigue after autologous bone marrow transplantation for breast cancer	14	Aetiology	
A psychosocial group intervention reduced psychological distress and enhanced coping in primary breast cancer	15	Children who participated in school extracurricular activities were less likely to drop out or to have been arrested	29
Review: advice from doctors and nurses, behavioural interventions, nicotine replacement treatment, and several pharmacological treatments increase smoking cessation rates	16	Genetic and specific environmental risk factors affected comorbidity of major depression and alcoholism	30
Lithium reduced aggression and was safe in aggressive children and adolescents with conduct disorder admitted to hospital	17	Qualitative	
Disulfiram was effective in patients with cocaine abuse who were also methadone treated opioid addicts	18	Psychiatric nurses expressed conflicting feelings in the decision to use physical restraints	31
In hospital cognitive rehabilitation was as effective as home rehabilitation in moderate to severe traumatic brain injury	19	Inpatients with personality disorders who stopped treatment had problems with the institutional culture and relationships	32
		Glossary	Inside back cover