

Evidence-Based Mental Health

Contents

Purpose and procedure	34	Cognitive behavioural therapy was more effective for panic disorder when given with 6 hours of total therapist contact than with 2 or 0 hours	53
Other articles noted	36	Review: SSRIs and tricyclic antidepressants increase response rates in post-traumatic stress disorder in the short term.....	54
EBMH notebook		Review: behavioural treatment has a small to moderate beneficial effect on chronic low back pain	55
Of studies, summaries, synopses, and systems: the “4S” evolution of services for finding current best evidence ..	37	Hypericum extract was as effective as imipramine but had fewer adverse effects in mild to moderate depression.....	56
Evidence-based psychiatric/mental health nursing	39		
Therapeutics		Prognosis	
The addition of folic acid to fluoxetine for major depression increased response rates especially in women	41	Poor insight, social avoidance, and better coping predicted prolonged duration of untreated psychosis in schizophrenia.....	57
Cognitive behavioural therapy improved most sleep outcomes in insomnia secondary to chronic pain.....	42	Review: women have worse outcomes than men after traumatic brain injury.....	58
Review: selective serotonin reuptake inhibitors reduce symptoms in premenstrual syndrome	43	Approximately two thirds of adolescents with newly diagnosed eating disorders recovered within 5–11.5 years	59
Review: no evidence exists that antidepressants acting at > 1 pharmacological site differ in efficacy from SSRIs in major depression	44	Aetiology	
Review: antidepressants plus benzodiazepines lead to fewer dropouts and less depression severity at 4 weeks in major depression.....	45	Depression was associated with increased risk of coronary heart disease and death in older adults	60
Review: behavioural interventions have a small to medium beneficial effect on diabetes management in adolescents with type 1 diabetes.....	46	Review: 14 risk factors for post-traumatic stress disorder include childhood abuse and family psychiatric history	61
No difference in effectiveness existed between community and hospital based interventions for children with behavioural disorders	47	Review: twin studies show that genes and individual environmental influences contribute to the aetiology of major depression	62
Review: evidence on anxiolytic drugs for smoking cessation is inconclusive.....	48	Quality improvement	
Nurse telehealth care reduced depressive symptoms and improved functioning	49	Pharmacist medication review decreased death, and number and cost of drugs prescribed for residents in nursing homes.....	63
A cognitive behavioural family intervention reduced psychiatric morbidity in caregivers of patients with Alzheimer’s disease	50	Economics	
Partner support reduced depressive symptoms in postpartum depression	51	Intensive case management was not more cost effective than standard case management for severe psychosis....	64
Virtual reality exposure was as effective as standard exposure for reducing fear of flying	52	Glossary	Inside back cover