

Evidence-Based Mental Health

Contents

Purpose and procedure	66	Guided threat focus and reappraisal was better than safety seeking behaviour for reducing fear in claustrophobia.....	86
EBMH notebook			
1. Enabling evidence-based change in health care	68	Intensive intervention for fear of childbirth did not reduce requests for caesarean section but decreased duration of vaginal labour	87
Other articles noted	72		
Therapeutics			
Review: sildenafil improves sexual and erectile function in men	73	Review: selegiline leads to a small short term improvement in cognition and activities of daily living in Alzheimer's disease.....	88
Review: antidepressants increase remission and clinical improvement in bulimia nervosa.....	74	The addition of olanzapine to valproate or lithium for acute manic or mixed bipolar episodes reduced manic symptoms	89
Review: psychological treatment is as effective as antidepressants for bulimia nervosa, but a combination is best	75	Extract of <i>Ginkgo biloba</i> added to haloperidol was effective for positive symptoms in refractory schizophrenia.....	92
A school based psychosocial intervention was effective in children with persistent post-disaster trauma symptoms	76	Diagnosis	
Risperidone reduced the risk of relapse in outpatient schizophrenia and schizoaffective disorder	77	2 clock tests had high accuracy and naïve raters had acceptable accuracy for detecting dementia	91
A group cognitive therapy programme reduced depressive episodes in at risk adolescents at 12 months of follow up.....	78	Aetiology	
A behavioural management for continence intervention reduced urinary incontinence symptoms in older rural women	79	Review: major anxiety disorders all have substantive familial aggregation.....	92
Naltrexone for 3 or 12 months in addition to psychosocial counselling did not reduce drinking in alcohol dependence	80	Mental disorders predicted all cause and cause specific deaths in Finnish adults.....	93
Review: methadone dose reduction is better than α_2 adrenergic agonists for treatment retention in opioid dependence	81	High trait anger increased stroke in people ≤ 60 years and those with high density lipoprotein cholesterol concentrations > 47 mmol/l.....	94
Review: antidepressants are not effective for cocaine dependence	82	Quality improvement	
Review: dopamine agonists are not effective for cocaine dependence	83	Case management reduced hospital admission in caregivers of people with Alzheimer's disease.....	95
A multidisciplinary community based rehabilitation programme improved social functioning in severe traumatic brain injury.....	84	Qualitative	
Review: amisulpride is effective and safe for schizophrenia.....	85	4 thematic patterns described the experience of placing a relative with Alzheimer's disease in a special care unit.....	96
		Glossary	Inside back cover