

Evidence-Based Mental Health

Contents

Purpose and procedure	98	Group psychoeducation reduces recurrence and hospital admission in people with bipolar disorder	115
EBMH Notebook		Collaborative care speeds recovery from depression.....	116
The Art of Evidence-Based Child Psychiatry	99	Review: selective serotonin reuptake inhibitors as effective as tricyclic antidepressants for major depression, and may have fewer adverse effects.....	117
Bipolar disorder.....	101	Review: repetitive transcranial magnetic stimulation is of unknown effectiveness in people with depression	118
Depression in children and adolescents	103	Gradual reduction of benzodiazepines, with or without cognitive behavioural therapy, increases successful withdrawal rates compared with no support in long-term users	119
Prevalence		Helping parents solve stressful life problems enhances the effectiveness of interventions for child aggression and antisocial behaviour	120
Prodromal symptoms may be identified by people with bipolar or unipolar depression.....	105	Review: evidence is lacking about suicide prevention in young people.....	121
Risk of suicide remains high fifteen years after deliberate self-harm.....	106	Review: therapeutic relationship is modestly correlated with treatment outcome in child and adolescent psychotherapy.....	122
Expected levels of behaviour problems in a population sample of children between four and eighteen years.....	107	Behavioural family counselling reduces drug use in opioid-dependent men	123
Diagnosis		Trauma-focused group psychotherapy is not effective for posttraumatic stress disorder in Vietnam veterans.....	124
Insufficient evidence on benefits of early detection and treatment of dementia	108	Self-help manuals may reduce bingeing and purging in bulimia nervosa	125
Nurse administered structured clinical interviews improve psychiatric diagnosis and treatment in community settings .	109	Review: stress is a problem for mental health nurses but research on interventions is insufficient	126
Therapeutics		Economics	
Rofecoxib or naproxen do not slow progression of mild to moderate Alzheimer's disease	110	Counselling may be more expensive than usual care for people with mental health problems, but may not be more effective for improving Beck depression score in the long term	127
Oestrogen plus progestin doubles the risk of dementia in post-menopausal women	111	Qualitative	
Targeted cognitive behavioural therapy may reduce relapse in people with prodromal symptoms of schizophrenia	112	Stigma alone does not explain non-disclosure of psychological symptoms in general practice.....	128
Review: the case for antidepressants for treating depression in people with schizophrenia remains unproven	113		
Family focused psychoeducational therapy decreases relapse and rehospitalisation in people with a manic episode and bipolar disorder	114		