

Evidence-Based Mental Health

Contents

EBMH Notebook

Schizophrenia 2

Prognosis

58% of people recovering from a major depressive episode are unlikely to relapse after 5 years..... 5

Modern treatments for severe depression still have poor long term outcome 6

Diagnosis

The minimum data set depression rating scale (MDSDRS) lacks reliability for identifying depression among older adults living in nursing homes 7

Inclusion of informant ratings of cognitive difficulties improves the accuracy of the MMSE in predicting Alzheimer's disease 8

The self rating inventory for post-traumatic stress disorder may aid diagnosis among older people in the community 9

Therapeutics

Sertraline improves symptoms in children and adolescents with major depressive disorder 10

Review: individual psychotherapy may be an effective treatment for people with bulimia nervosa..... 11

Flexible oral olanzapine dosing in acutely agitated people is immediately effective in reducing symptoms 12

Aripiprazole is effective against the symptoms of schizophrenia and schizoaffective disorder 13

Review: new generation antipsychotic drugs may reduce relapse rates more effectively than conventional antipsychotic drugs in people with schizophrenia..... 14

Manual assisted cognitive behaviour therapy is as effective as treatment as usual for deliberate self harm, but is more cost effective 15

Review: psychodynamic therapy and cognitive behavioural therapy are effective in the treatment of personality disorders 16

High dose pregabalin is effective for the treatment of generalised anxiety disorder 17

Combined imaginal exposure and cognitive restructuring therapy is more effective than supportive counselling for treating post-traumatic stress disorder 18

Cognitive stimulation therapy improves cognition and quality of life in older people with dementia..... 19

Donepezil slows decline in daily life activities in people with moderate to severe Alzheimer's disease and alleviates caregiver burden..... 20

Outreach supported antidepressant treatment and cognitive behavioural therapy are effective for depression in low income minority women 21

Treating low perceived social support and depression after myocardial infarction does not increase event-free survival 22

Supervised co-prescription of heroin to treatment-resistant heroin addicts is more effective than treatment with methadone alone 23

Aetiology

Adolescent major depressive disorder may predict poor psychosocial functioning in young adulthood..... 24

Borderline personality disorder, drug use disorder, and worsening depression or substance abuse are significant predictors of suicide attempts in people with Axis I and II disorders 25

Mini-clusters of potentially prodromal symptoms may identify psychiatrically well Amish children at higher risk of developing bipolar I disorder..... 26

Review: at least one third of people report persistent memory loss after electroconvulsive therapy 27

Review: exploration of psychological and physical health differences between caregivers and non-caregivers 28

Review: psychodynamic therapy and cognitive behavioural therapy are effective in the treatment of personality disorders 16

Miscellaneous

Other articles noted 3