

Evidence-Based Mental Health

Contents

Purpose and procedure	94	Withdrawing neuroleptic treatment from people with dementia, does not result in a significant change to their behaviour, psychiatric symptoms, or quality of life	111
EBMH notebook			
Evidence-based medicine: ethically obligatory or ethically suspect?.....	96	Donepezil did not reduce the rate of institutionalisation or disability in people with mild to moderate Alzheimer's disease	112
Response to Dr Gupta.....	97	Trauma focused cognitive behavioural therapy reduces PTSD more effectively than child centred therapy in children who have been sexually abused.....	113
Editorial			
The worm turns: publication bias and trial registers revisited	98	Switching serotonin reuptake inhibitors may be of benefit in people with obsessive compulsive disorder	114
Prognosis			
Duration of untreated psychosis significantly associated with positive symptoms one year after treatment.....	101	Review: relying on published data alone overestimates efficacy and safety of SSRIs in children.....	115
Acute stress disorder is of limited benefit in predicting post-traumatic stress disorder in people surviving traumatic injury	102	Increasing a partner's understanding of motherhood significantly reduces postnatal distress and depression in first time mothers with low self esteem.....	116
Psychological function is mildly impaired for prolonged periods before and after first unipolar major depressive episode.....	103	Integrative Behavioral Couple Therapy is as effective as Traditional Behavioral Couple Therapy for distressed couples.....	117
Therapeutics			
Review: in people with schizophrenia, lithium is ineffective as sole therapy, while evidence on augmenting antipsychotics with lithium is inconclusive	104	Cognitive behavioural therapy significantly improves hypochondriacal symptoms	118
Cognitive behavioural therapy improves psychotic symptoms at 18 months in people with schizophrenia	105	Aetiology	
Review: high doses of antipsychotic drugs are no more or less effective than medium doses in people with schizophrenia.....	106	Review: current evidence does not show a strong causal relation between the use of cannabis in young people and psychosocial harm.....	119
Review: combining pindolol with an SSRI improves early outcomes in people with depression.....	107	Concomitant loop diuretics and ACE inhibitors increase risk of lithium toxicity in elderly people	120
Review: antidepressants are moderately effective in people with depression and a substance dependence disorder.....	108	Externalising problems and environmental stressors predict adolescent problems in children from divorced families	121
Review: St John's Wort may be less effective than previously thought in people with depression.....	109	Individuals with opioid dependence who leave drug treatment have higher overdose and death rates than those still in treatment.....	122
PEARLS home based treatment significantly improves depression, dysthymia, and health related quality of life in older people	110	Prevalence	
Miscellaneous			
		Depressive disorders were the fourth leading cause of global disease burden in the year 2000.....	123
		Personality disorders commonly occur in people with alcohol and drug use disorders	124
		Other articles noted	100