

Evidence-Based Mental Health

Contents

Purpose and procedure	90	Collaborative care depression management for older adults: level of comorbidity does not affect outcome ..	105
EBMH notebook		Collaborative care management improves physical functioning in older people with depression	106
Risk assessment	91	St John's wort is at least as effective as paroxetine in reducing severity of depression and is better tolerated	107
With apologies to Albert: everything is <i>not</i> relative	93	Postnatal home visits from healthcare professionals show promise for preventing postnatal depression.....	108
Other articles noted	94	Adding hypnosis to cognitive behavioural therapy may reduce some acute stress disorder symptoms	109
Prognosis		Coordinated care consisting of cognitive behavioural therapy plus medication improves panic disorder.....	110
The incidence of delirium in older people with a mood disorder is similar with lithium and valproate	95	Pharmacotherapy is an effective treatment option for generalised anxiety disorder	111
Lifetime risk of suicide in people with schizophrenia lower than commonly reported	96	Methadone plus contingency management or performance feedback reduces cocaine and opiate use in people with drug addiction	112
Attendance at accident and emergency for deliberate self harm predicts increased risk of suicide, especially in women.....	97	Aetiology	
Early onset of drinking increases alcohol use in adulthood.....	98	Forensic database study suggests selective serotonin reuptake inhibitors do not increase the risk of suicide in people taking antidepressants	113
Diagnosis		Suicide risk peaks in first week of psychiatric hospitalisation and post-discharge.....	114
The sentence completion test for depression can distinguish between people with and without major depressive disorder	99	Atypical antipsychotic use during the first trimester of pregnancy may not increase major malformations.....	115
Therapeutics		Prevalence	
Long acting injectable naltrexone is effective and safe for treating alcohol dependence	100	Comorbid pain related somatisation is common in Latin Americans with major depressive disorder	116
Bipolar disorder in young people: divalproex sodium no more effective than lithium for maintenance	101	Prescription of pharmacotherapy for depression in elderly people varies with age, race, gender, and length of care	117
Adding cognitive therapy to minimal psychiatric care prevents short term, but not long term, relapse in people with bipolar disorder	102	Fifteen per cent of people treated for mental health disorders are homeless	118
Cognitive behavioural therapy reduces psychological distress in carers of people with Parkinson's disease ..	103		
Some psychosocial therapies may reduce depression, aggression, or apathy in people with dementia	104		