

Evidence-Based Mental Health

Contents

Purpose and procedure	2	Pharmacotherapy and cognitive behavioural therapy: similarly cost effective compared with community referral for disadvantaged women with major depression	16
Other articles noted	4	A structured activity programme reduces depressive symptoms in moderately depressed older men with coronary heart disease, but not women	17
Prognosis		Training nursing home care staff to recognise psychopathology improves their ability to identify depressed residents.....	18
Poor social and interpersonal functioning prior to diagnosis predicts poor outcome for people with first episode psychosis.....	5	Review: cholinesterase inhibitors reduce burden and care time for informal carers of people with Alzheimer's disease.....	19
Therapeutics		Neither vitamin E nor donepezil delays progression from amnesic mild cognitive impairment to Alzheimer's disease in the long term	20
Risperidone improves restricted, repetitive, and stereotyped behaviour in autistic children and adolescents	6	Review: light therapy is an effective treatment for seasonal affective disorder	21
Atomoxetine improves teacher rated symptoms in children with ADHD more than placebo	7	Supported employment interventions are effective for people with severe mental illness.....	22
Multisystemic therapy reduces long term rearrest compared with usual treatment	8	Aetiology	
Intramuscular haloperidol and olanzapine begin to reduce psychosis within 24 hours	9	Epilepsy or a family history of epilepsy increases the risk of schizophrenia or schizophrenia-like psychosis.....	23
Review: no evidence to support gradual over abrupt switching of antipsychotics	10	History of psychiatric hospital admission increases the risk of suicide less in the very old than in middle aged people.....	24
Individual dietary education reduces olanzapine associated weight gain	11	Prevalence	
Cognitive behavioural therapy does not improve outcome in obese women with binge eating disorder receiving a comprehensive very low calorie diet programme	12	Suicide ideation and attempts are more prevalent in people aged 25–44 years in Australia but become less prevalent in older people.....	25
Cognitive behavioural therapy reduces nocturnal panic in people with panic disorder.....	13	Young people admitted with self-injury had more severe psychiatric disorders in 2000 than in 1990.....	26
Cognitive therapy is more cost effective than standard care alone for bipolar disorder	14	About 26% of people in the US have an anxiety, mood, impulse control, or substance disorder	27
Divalproex and lithium are similarly cost effective for adults with bipolar disorder	15	About 30% of men with schizophrenia or schizoaffective disorders have obsessive-compulsive symptoms.....	28