

Evidence-Based Mental Health

Contents

Purpose and procedure	30	First episode psychosis: olanzapine and haloperidol provide similar improvements in quality of life and social functioning	47
EBMH notebook			
APA, NAMI, NMHA, and evidence-based behavioural medicine: a comment	32	Adding sarcosine, but not D-serine, to risperidone improves symptoms in people with acute phase schizophrenia	48
Other articles noted	35	Telephone delivered cognitive behavioural therapy improves depression in people with multiple sclerosis faster than telephone delivered supportive emotion-focused therapy	49
Prognosis			
Review: longer duration of untreated psychosis is associated with worse outcome in people with first episode psychosis	36	Counselling and cognitive behavioural therapy reduce anxiety and depression in women with postnatal depression	50
Cognitive function is not a prognostic factor for long term functional status in older people	37	A universally administered primary prevention programme for depression reduces symptoms in German adolescents with low self-efficacy	51
Diagnosis			
The short form adult attention deficit/hyperactivity self-report scale is a useful diagnostic measure	38	Continued escitalopram reduces risk of relapse in people with generalised social anxiety disorder	52
Adding a question that asks whether help is wanted improves sensitivity of a depression screening tool	39	Combination treatment of behaviour therapy and medication significantly reduces obsessive-compulsive symptoms in people who have responded to pharmacotherapy	53
Therapeutics			
Risperidone effectively increases rates of symptomatic remission of acute mania in people with bipolar disorder	40	Review: atypical antipsychotic drugs slightly increase mortality rates in elderly people with dementia	54
Aripiprazole reduces acute mania compared with haloperidol in bipolar I disorder and causes fewer adverse effects	41	Abstinence incentives improve stimulant abusers' compliance with community-based psychosocial treatments	55
Olanzapine plus fluoxetine reduce depressive symptoms faster than either drug alone in people with treatment resistant depression	42	Discontinuing risperidone results in relapse in children with autism spectrum disorders	56
Review: cognitive behavioural therapy is more effective than control and similarly effective to pharmacotherapy for generalised anxiety disorder	43	Aetiology	
Review: psychological treatments improve pathological gambling in the short and long term	44	Adolescent panic attacks are associated with increased risk of personality disorder as a young adult	57
Methylphenidate hydrochloride reduces hyperactivity in children with pervasive development disorders	45	Anxiety disorders are a risk factor for suicidal behaviour	58
Integrated care improves one year outcomes in first episode psychosis	46	Prevalence	
		Major depressive disorder is common in the US among Native Americans, women, the middle aged, the poor, and widowed, separated, or divorced people	59
		Miscellanea	
		Correction	34