

Evidence-Based Mental Health

Contents

Purpose and procedure	62	IMPACT collaborative care improves depression in elderly patients in primary care in the longer term....	76
Other articles noted	64	Vardenafil improves comorbid erectile dysfunction and mild depression	77
Prognosis		Review: an early sustained response can occur in the first and second weeks of antidepressant treatment for major depressive disorder.....	78
Early discontinuation of antidepressant treatment in adults is very common in the USA	65	Review: case management in primary health care improves symptoms and drug adherence in people with major depression	79
Diagnosis		Review: cognitive behavioural therapy is an effective treatment for depression, panic disorder, and generalised anxiety disorder, but may be less effective in severe cases.....	80
The Anxiety Change Expectancy Scale score shows high internal validity and correlation with validated measures of anxiety, self-esteem, and hopelessness in varied settings.....	66	Case management and vouchers improve uptake of methadone treatment programmes.....	81
The clinician rated CY-BOCS scale is more sensitive than the self-reported LOI-CV in children and adolescents with obsessive compulsive disorder.....	67	Aetiology	
Therapeutics		No apparent difference in suicide risk between older and newer antidepressants although older drugs may increase risk of suicide attempt during the first month of treatment	82
Modafinil improves symptoms of ADHD compared with placebo in young people	68	Substance use and a history of psychosis are risk factors for serious adverse events in people with bipolar disorder.....	83
Early intervention has no effect on symptoms in people with first episode, non-affective psychosis, although it may improve overall function and medication adherence.....	69	Residual mood symptoms and number of previous episodes predict recurrence of bipolar disorder.....	84
Review: higher rates of attrition in antipsychotic treatment arms of placebo controlled trials than in trials with active comparators.....	70	Excessive alcohol consumption is not a risk factor for anxiety and depression, nor are anxiety and depression a risk factor for excessive alcohol consumption.....	85
Olanzapine and risperidone may improve neurocognition more than haloperidol in people with schizophrenia who continue treatment for 52 weeks ..	71	Prevalence	
Olanzapine is associated with more rapid weight gain than haloperidol in people with first episode psychosis	72	Lifetime illicit drug use and drug dependence are common by the age of 25 in New Zealand.....	86
Maintenance treatment with olanzapine reduces relapse in people with bipolar I disorder who have responded to acute olanzapine treatment	73	Review: high rates of substance abuse or dependence in male and female prisoners.....	87
Topiramate improves psychopathological symptoms and quality of life in women with borderline personality disorder.....	74	Lifetime prevalence of social anxiety disorder in the USA is 5%	88
Mirtazapine reduces social anxiety and improves quality of life in women with social phobia.....	75		