

Evidence-Based Mental Health

Contents

| | | | |
|---|-----|---|-----|
| Purpose and procedure | 90 | Donepezil improves cognitive function and reduces decline in activities of daily living in people with severe Alzheimer's disease | 104 |
| Other articles noted | 92 | Risperidone maintenance treatment increases time to symptom recurrence in young people with disruptive behaviour disorders | 105 |
| Prognosis | | Schizophrenia: risperidone and olanzapine increase time to discontinuation compared with quetiapine and ziprasidone | 106 |
| Long term risk of death in people who have self-harmed is higher than in the general population | 93 | Aetiology | |
| One in 10 elderly people with minor or subsyndromal depression develops major depression within a year... | 94 | Gender, age, ethnicity and area of residence influence incidence of psychotic disorders | 107 |
| High rates of recurrence of depression in children and adolescents | 95 | High risk cognitive style predicts onset of depression | 108 |
| Therapeutics | | Entering menopause increases the risk of first episode depression | 109 |
| Universal and targeted cognitive behavioural approaches do not reduce depression in young people at risk of depression | 96 | The menopausal transition increases the risk of depressive symptoms and depression diagnosis in women without a history of depression | 110 |
| The Youth-Nominated Support Team for Suicidal Adolescents (version 1) intervention does not reduce suicidal attempts | 97 | Depressive symptoms in elderly people are associated with an increased risk of developing mild cognitive impairment independently of vascular disease | 111 |
| Review: psychotherapy plus antidepressant therapy increases response rate in people with panic disorder more than either treatment alone | 98 | Among middle aged men in Japan, non-drinkers and regular heavy drinkers are more likely to commit suicide than occasional drinkers | 112 |
| Cognitive behavioural therapy does not reduce overall episode recurrence in people with recurrent bipolar disorder | 99 | Prevalence | |
| Major depressive disorder: similar remission rates with bupropion, sertraline, or venlafaxine following treatment switch from citalopram | 100 | Lifetime mood, anxiety, and drug use disorders are common in the United States population | 113 |
| Continuing paroxetine treatment reduces recurrence of major depression in the elderly | 101 | Lifetime prevalence of panic disorder is about 5% in the USA | 114 |
| Educational workshops and decision support software increase detection of dementia in the elderly in primary care settings | 102 | Review: worldwide lifetime prevalence of anxiety disorders is 16.6%, with considerable heterogeneity between studies | 115 |
| Training nursing home staff to improve psychosocial care reduces neuroleptic use in people with dementia without an increase in agitation | 103 | Adult attention deficit hyperactivity disorder is common among people in the US who are male, white, previously married, or not employed | 116 |