

Evidence-Based Mental Health

Contents

Purpose and procedure	90	Donepezil improves cognitive function and reduces decline in activities of daily living in people with severe Alzheimer's disease	104
Other articles noted	92	Risperidone maintenance treatment increases time to symptom recurrence in young people with disruptive behaviour disorders	105
Prognosis		Schizophrenia: risperidone and olanzapine increase time to discontinuation compared with quetiapine and ziprasidone	106
Long term risk of death in people who have self-harmed is higher than in the general population	93	Aetiology	
One in 10 elderly people with minor or subsyndromal depression develops major depression within a year...	94	Gender, age, ethnicity and area of residence influence incidence of psychotic disorders	107
High rates of recurrence of depression in children and adolescents	95	High risk cognitive style predicts onset of depression	108
Therapeutics		Entering menopause increases the risk of first episode depression	109
Universal and targeted cognitive behavioural approaches do not reduce depression in young people at risk of depression	96	The menopausal transition increases the risk of depressive symptoms and depression diagnosis in women without a history of depression	110
The Youth-Nominated Support Team for Suicidal Adolescents (version 1) intervention does not reduce suicidal attempts	97	Depressive symptoms in elderly people are associated with an increased risk of developing mild cognitive impairment independently of vascular disease	111
Review: psychotherapy plus antidepressant therapy increases response rate in people with panic disorder more than either treatment alone	98	Among middle aged men in Japan, non-drinkers and regular heavy drinkers are more likely to commit suicide than occasional drinkers	112
Cognitive behavioural therapy does not reduce overall episode recurrence in people with recurrent bipolar disorder	99	Prevalence	
Major depressive disorder: similar remission rates with bupropion, sertraline, or venlafaxine following treatment switch from citalopram	100	Lifetime mood, anxiety, and drug use disorders are common in the United States population	113
Continuing paroxetine treatment reduces recurrence of major depression in the elderly	101	Lifetime prevalence of panic disorder is about 5% in the USA	114
Educational workshops and decision support software increase detection of dementia in the elderly in primary care settings	102	Review: worldwide lifetime prevalence of anxiety disorders is 16.6%, with considerable heterogeneity between studies	115
Training nursing home staff to improve psychosocial care reduces neuroleptic use in people with dementia without an increase in agitation	103	Adult attention deficit hyperactivity disorder is common among people in the US who are male, white, previously married, or not employed	116