

Adjusted odds ratios for suicide by phase of admission or discharge and by number of hospitalisations

	<b>OR for women (95% CI)*</b>	<b>OR for men (95% CI)*</b>
<b>Admission period</b>		
First week	81.6 (52.1 to 127.7)	59.8 (42.1 to 85.1)
Weeks 2 to 4	71.0 (49.9 to 101.0)	51.9 (38.2 to 70.7)
1–6 months	49.8 (37.9 to 65.6)	30.4 (24.1 to 38.3)
7–12 months	37.2 (19.8 to 69.9)	9.9 (6.2 to 15.7)
1 year and over	1.5 (0.8 to 3.0)	1.6 (1.1 to 2.6)
<b>Discharge period</b>		
First week	246.0 (152.5 to 396.6)	102.0 (73.2 to 142.1)
Weeks 2–4	90.7 (66.0 to 124.6)	62.1 (48.4 to 79.6)
1–6 months	35.8 (29.1 to 44.0)	28.4 (24.0 to 33.5)
7–12 months	19.3 (15.6 to 23.9)	14.7 (12.3 to 17.7)
1–3 years	10.5 (8.7 to 12.6)	6.4 (5.5 to 7.4)
4–5 years	5.4 (4.4 to 6.6)	4.6 (3.9 to 5.5)
Over 5 years	3.0 (2.6 to 3.6)	2.8 (2.5 to 3.3)

\*Comparisons versus people with no admission history.