

Webextra Table: Essential and potentially inappropriate use of antipsychotics by income

	Income Quintile OR (95% CI)				
	Highest (ref)	Fourth	Third	Second	Lowest
Adults with schizophrenia (essential use)					
Antipsychotic use (any)	1.00	1.22 (1.04 to 1.44)	1.71 (1.44 to 2.04)	1.08 (0.92 to 1.26)	1.05 (0.90 to 1.23)
Long-term essential antipsychotic use	1.00	1.31 (1.15 to 1.48)	1.87 (1.64 to 2.14)	1.40 (1.23 to 1.59)	1.18 (1.04 to 1.34)
Older adults with dementia (potentially inappropriate use)					
Antipsychotic use (community dwelling adults)	1.00	1.05 (0.93 to 1.19)	1.05 (0.93 to 1.19)	1.19 (1.05 to 1.34)	1.24 (1.10 to 1.39)
Antipsychotic use (adults in long-term care)	1.00	1.02 (0.89 to 1.18)	0.94 (0.81 to 1.08)	0.88 (0.76 to 1.02)	0.90 (0.78 to 1.04)
Long-term antipsychotic use (community dwelling adults)	1.00	0.97 (0.84 to 1.12)	0.93 (0.80 to 1.08)	1.10 (0.96 to 1.27)	1.19 (1.03 to 1.36)
Long-term antipsychotic use (adults in long-term care)	1.00	1.05 (0.91 to 1.21)	1.05 (0.91 to 1.21)	0.95 (0.82 to 1.09)	0.96 (0.83 to 1.10)