

WebExtra Table 1: Lifetime prevalence (%) of DSM-IV or CIDI-assessed mental health disorders amongst adolescents with and without lifetime suicidality

Disorder	Total sample			Lifetime ideators		No suicidal behaviours (n=5766)
	Ideation (n=717)	Plan (n=203)	Attempt (n=196)	Planned attempt (n=112)	No plan attempt (n=84)	
Fear and anger disorders						
Specific phobia	36.8*	39.0*	44.3*	40.7*	49.6*	17.6
PD/A	10.1*	10.6*	10.4	10.8	9.8	4.0
Social phobia	19.9*	16.2*	25.6*	17.7*	37.5	7.0
IED	29.4*	35.7*	35.2*	42.1*	24.9	11.5
Distress disorders						
SAD	11.9	8.9	7.0	3.9	11.5	7.0
PTSD	16.1*	27.9*	25.7*	34.2*	13.0	3.1
MDD/DYS	56.8*	69.7*	75.7*	76.7*	74.3*	13.3
GAD	8.4*	10.3	9.2*	11.2	6.2	1.6
Disruptive behaviour disorders						
ADHD	16.3*	19.0*	21.5*	23.5	18.5	7.0
ODD	34.4*	41.6*	50.0*	51.5*	47.8*	9.6
Conduct disorder	20.2*	22.8*	33.5*	26.9*	43.3*	5.0
Eating disorders	15.8*	11.9	26.7*	15.3	43.7*	4.0
Substance abuse						
Alcohol abuse	18.4*	28.6*	24.3*	34.4*	9.2	4.6
Illicit drug abuse	27.4*	28.0*	34.7*	27.5*	45.3*	6.4
Biopolar I or II	9.1*	11.9*	13.2*	18.6*	5.2	2.2
Any disorder	89.3*	93.6*	96.1*	96.7*	95.2*	49.5
* Significant difference in prevalence from adolescents with no history of suicidal behaviour; p<0.05						